

**JUCD Adult Covid-19 (C19) Vaccine Allergy Service Pathway:
Managing Patient with a History of Allergic Reactions
For GPs**

**Table 1: First Dose Advice for Managing patients with history of Allergy or Anaphylaxis
who have NOT yet received C19 vaccine**

PATIENT CHARACTERISTICS

Previous allergic reaction (including anaphylaxis¹) to a food, insect sting and most medicines (where trigger has been identified)

Family history of allergies
Previous non-systemic reaction to a vaccine

Hypersensitivity to non-steroidal anti-inflammatory drugs e.g. aspirin, ibuprofen

Chronic Spontaneous Urticaria or Mastocytosis



Proceed with vaccination as normal. According to local guidelines.

PATIENT CHARACTERISTICS

Confirmed diagnosis of PEG allergy or reaction to macrogol



GP to complete electronic referral to:

CoViD Vaccine Allergy-ADVICE AND GUIDANCE ONLY-Royal Derby - UHDB-RTG - Service ID 7974261

PATIENT CHARACTERISTICS

History of immediate anaphylaxis¹ to multiple drug classes to non-COVID vaccine or biologics e.g. Infliximab, adalimumab, tocilizumab, secukinumab, ustekinumab, anakinra.


History of unexplained anaphylaxis



GP to complete electronic referral to:

CoViD Vaccine Allergy-ADVICE AND GUIDANCE ONLY-Royal Derby -UHDB-RTG - Service ID 7974261

Flowchart for managing patients who have allergic reactions to a previous dose of COVID-19 vaccine



1 Consider pre-treatment with non-sedating antihistamine, at least 30mins prior to vaccination.