

The Derbyshire Nutrition and Hydration Pack for Care Homes

This resource pack has been designed to promote excellence in nutrition and hydration care in the care home setting.

If you have any queries about the information within this booklet, please contact:

- The North Derbyshire Dietitians on: 01246 512173
- The South Derbyshire Dietitians on: 01332 258214

Endorsed by:



Contents	
Page 2	Introduction
Page 2	Nutritional Screening - MUST
Page 6	End of Life Care
Page 6	Dysphagia & Texture Modified Diets
Page 7	Promoting a Positive Meal Experience
Page 8	Healthy Eating
Page 9	Vegetarian & Vegan Diets
Page 10	Food Fortification
Page 12	Vegan Food Fortification
Page 13	Nourishing Snacks
Page 13	Nourishing Snack Ideas
Page 14	Nourishing Drinks
Page 17	Nourishing Puddings
Page 18	Example Fortified Menu
Page 19	Eating Well With Dementia
Page 21	Diabetes
Page 24	Vitamin D
Page 25	Coeliac Disease
Page 26	Hydration
Page 27	Food Charts
Page 28	Over The Counter Nutrition Supplements
Page 28	Oral Nutrition Supplements Which Can Be Prescribed
Page 29	Referral to Nutrition & Dietetic Services
Page 30	Resource Section
Page 31	Monthly Catering Communication Sheet
Page 33	The Eatwell Guide
Page 34	Algorithm for Nutrition Care Planning
Page 35	Texture Modified Diet, Fluids & IDDSI
Page 37	BAPEN MUST Tools
Page 41	Useful Web Links

Introduction

The Nutrition and Dietetic Services have produced this guide to help nursing and residential care homes in your locality to support staff to provide all residents with excellent nutritional care, in line with Care Quality Commission (CQC) Regulation 14: Meeting Nutritional And Hydration Needs. It is essential that all care homes must meet Regulation 14 by providing nutrition and hydration appropriate to their residents' individual nutritional needs to be an appropriate care provider.

The care home population is diverse with a wide range of nutrition and hydration needs and challenges. Those in residential care settings may have a requirement for 'healthy eating' or 'eating for health'. 'Eating for health' better reflects the needs of residents who require a special / therapeutic / more energy dense diet. Each resident requires a diet that meets their own health and nutrition needs. Good nutrition and hydration is fundamental in health and disease and impacts on recovery times, quality of life and health outcomes including premature death.

Nutritional Screening

All residents admitted to care homes should be screened for malnutrition using an accredited screening tool, such as the Malnutrition Universal Screening Tool (MUST) produced by BAPEN (see page 37 – 40, link also in the Useful Web Links section, page 41).

Nutritional screening should be completed **within 24 - 48 hours** of each resident's admission.
and screening should be **re-assessed at least monthly** for all residents.

The MUST screening consists of 5 steps:

- Step 1: BMI score
- Step 2: Weight loss score
- Step 3: Acute disease effect score
- Step 4: Overall risk of malnutrition
- Step 5: Management guidelines

Some care homes will have their own nutritional screening tool and if this is the case the dates along with the weight, heights and scores should be documented on this tool.

Screening Step 1: BMI Score

The Body Mass Index (BMI) should use a resident's **most recent weight**. Weight should be recorded in kilograms (kg) and the resident's height should be recorded in metres (m).

To calculate the BMI score, a BMI chart can be used (see page 38) or the following equation:

$$\text{Body mass index} = \frac{\text{weight (kg)}}{\text{height (m)}^2}$$

Record the BMI in the resident's notes, and the MUST score based on the information in the table below.

BMI	BMI MUST score	Nutritional risk
Greater than 20	0	Low
Between 18.5 and 20	1	Medium
Less than 18.5	2	High

Guidance on obtaining a weight:

- Measure weight at least monthly for all residents using suitable, regularly calibrated standing, seated, hoist or wheelchair scales.
- If your resident's weight varies greatly from the last documented weight, re-weigh them to ensure that no errors have occurred.
- To promote accuracy, aim to record resident's weights at a similar time each month, at a similar time of day, and in light clothing. Ensure that the floor surface is level.
- If you are unable to weigh a resident, you can instead estimate BMI using mid upper arm circumference (MUAC) (see page 35). Please note that this estimated BMI should not be used to calculate the residents MUST score.

Adjusting Weight For Amputations

You may have residents with amputations and as a result, their BMI from their amputation weight is inaccurate. To calculate their adjusted weight, please use the following:

Below knee	Current weight (kg) x 1.063
Full leg	Current weight (kg) x 1.18
Forearm	Current weight (kg) x 1.022
Full arm	Amputation Calculation Current weight (kg) x 1.05

Guidance on obtaining a height measure:

- A standing height measurement in metres should be taken where possible on admission to a care home using either a stadiometer or tape measure secured to a wall. Height measurements do not need to be repeated unless there are concerns over the accuracy of the first measurement.
- If you can't obtain a resident's height, you can try to obtain an ulna length which will give you an estimated height (see page 40 for conversion chart). If this is not possible, you may be able to obtain a self-reported height or a height reported from a family member. If an alternative, estimated or reported height is used for calculating the MUST score this should be clearly documented. Alternatively, a height measure could also be obtained from the residents general practice records.



Top Tip: Set a weighing day for each section of your care home, to ensure that residents are weighed at a similar time each month e.g. the afternoon of the 1st day of each month.

Screening Step 2: Weight Loss Score

The weight loss score is calculated as a percentage, using the **most recent weight** and the resident's normal / usual weight **from the last 3-6 months**. Record the residents normal / usual weight on the MUST tool. It is common for your resident's weight to fluctuate within a small margin above or below their normal weight (+/- 5%).

Obtaining a normal / usual weight may be difficult for a new resident. This might be available from the resident's general practice records, from the resident themselves or their relatives. If the resident's weight history is not available over the past 3-6 months, you will be unable to complete this step. Instead, complete Step 1 and Step 3 and record the reason why Step 2 has not been completed.

To determine percentage weight loss, you can use a chart (page 39 or a local version) or you can calculate it using the following equation (**weight must be in kg**):

$$\% \text{ weight loss} = (\text{previous weight} - \text{current weight}) \div \text{previous weight} \times 100$$

Record the % weight loss in the residents notes, and the MUST score for Step 2.

BMI	BMI MUST score	Nutritional risk
Less than 5%	0	Low
Between 5 – 10%	1	Medium
Greater than 10%	2	High



Top Tip: If you are unable to obtain a weight history for the resident or are finding it difficult to judge the weight loss, you could ask family, check with the GP, or use family photographs of the resident to visualise whether the resident has lost a significant amount of weight.

Screening Step 3: Acute Disease Effect

Step 3 generates a MUST score if a resident has been acutely **and** critically ill **and** there has been (or is likely to be) **NO** nutritional intake for more than 5 days. If these criteria are met, the resident would be given a MUST score of 2.

In most circumstances in the community setting, Step 3 **will not** apply to your residents. Therefore, for this step, most residents will be given a MUST score of 0.

Screening Step 4: Calculate Overall Risk of Malnutrition

To determine the overall MUST score, add the scores from Step 1, Step 2 and Step 3. Please see the below table:

Total MUST Score	Risk Of Malnutrition
0	Low
1	Medium
≥2	High

In a care home, the highest possible MUST score is usually 4. In *very rare circumstances* when a resident scores 2 for Step 3, the highest overall MUST score would be 6.

Screening Step 5: Management Guidelines and Care Planning

When you have determined the overall MUST score, refer to the Algorithm for Nutrition Care Planning to identify the appropriate care pathway for your resident's needs (see page 34).

Your resident's diet may need to change depending on their MUST score. Nutrition care plans and the aim of any nutritional interventions should be reviewed at least monthly. This is to reflect changing MUST scores and any changes to the management of their nutrition and hydration needs.

Regular communication with the catering staff will ensure that they are aware of any nutritional changes, and that all residents receive the appropriate diet for their needs. To document resident's nutritional needs conveniently for catering staff, the Monthly Catering Communication Sheet (see page 31-32) can be used.



Top Tip: Provide your general practice with a list of the resident's weights regularly (at least 3 monthly), so that they can update their medical records. This will improve the overall quality of integrated care provided.

End of Life Care

For residents who are in the last days of their life, their nutritional intake should maximise their quality of life by providing comfort, symptom relief and enjoyment of food. Nutritional screening, weighing the resident and prescriptions for oral nutritional supplements (for weight gain or maintenance) are **no longer appropriate** because they are not likely to provide any significant nutritional benefits.

During this time, the main nutritional priority for residents is to offer diet and fluids that are enjoyed and tolerated. This promotes comfort and quality of remaining life.

Dysphagia and Texture Modified Diets

Residents with difficulty in swallowing foods safely (oropharyngeal dysphagia) may require a texture modified diet. To reduce their risk of choking or aspiration (food or liquid entering their airway), a Speech and Language Therapist (SLT) may have suggested texture modifications to their foods and/or liquids. The British Dietetic Association (BDA) and the Royal College of Speech & Language Therapists (RCSLT) agreed to adopt the International Dysphagia Diet Standards Initiative (IDDSI) across the country from April 2019. For more information on IDDSI and its criteria, please see pages 35-36 or the link in the ‘Useful Web Links’ section.

A texture modified diet should only be given to residents following the advice of a healthcare professional, such as a SLT. Resident's need for a modified texture diet should be regularly reviewed, as their requirements for this may change over time. Changes should be communicated to **all** staff involved in a resident's food preparation and provision as well as family who may bring in snacks. It is important for texture modified meals to remain appetising and varied and balanced to meet the nutritional requirements of the residents. Every effort should be made to maintain the usual colour, nutrient content, flavour and appearance of foods served. Moulds can be used to improve the appearance of texture modified meals. Well planned menu rotation is also encouraged, to ensure that a variety of colourful, flavourful, appealing meals are served and to prevent taste fatigue for these residents as this may lead to a reduced food intake.

To achieve an appropriate texture modified menu as suggested by SLT, additional fluids are often added to foods to moisten or aid the pureeing process. This results in a larger volume of food which a resident is unlikely to eat and a dilution of the nutrients. To ensure that resident's receive adequate nutrition, texture modified meals often require fortification (please see page 10-18). If there is any concern regarding a residents' texture modified diet requirements, please contact their discharging SLT or Dietitian for more information.

Promoting a Positive Meal Experience

Before and during mealtimes, steps should be taken to promote a positive meal experience. The tips below will ensure your residents have a good meal experience and this can contribute to them eating more at meal times.

- Prepare residents for mealtimes.
- Encourage them to take eat their meals away from their beds and ideally at a dining table.
- Make them aware in advance about the menu options.
- Encourage residents or ensure that care staff wipe down the dining room tables or bed side tables before meals are served.
- Ensure all residents have an opportunity to visit the toilet and wash or wipe their hands before meals are served.
- Ensure that residents have the right equipment available at meal times e.g. glasses, hearing aids, well-fitting dentures, special cups, cutlery and plates if needed.
- Ensure all residents are sitting in a supported, upright position (45 degrees minimum).
- Keep the eating environment as calm as possible with minimal distractions. Some residents may eat better in company in a dining room whilst others may be better sitting alone in a lounge or in their own room. Consider their individual needs.
- Ask residents about their food and drink preferences at meal times.
- Aim for maximum independence during meal times but continue to observe, encourage and assist as needed.
- Help residents to open any packaging or cut up meals if needed.
- Ensure that all residents have their meal, suitable for their needs, as well as a suitable drink in reach.
- Avoid using large spout beakers. These deliver more fluid to a patient than they would normally swallow.
- Avoid all spouts if possible. If one is needed for a resident, use a small one.

Having to be fed can be emotive for many residents as they view this as further deterioration in their abilities. It is important to ensure that dignity is maintained when helping a resident to eat their meal. Implementing the following can help boost nutritional intake and again improve the residents' meal experience:

- When feeding a resident, sit slightly to one side at eye level.
- Feed at a pace appropriate to the resident, **do not** rush.
- Give the resident plenty of time to swallow.
- Tell the resident what you are feeding them at each mouthful and don't mix foods.
- Don't overload the spoon or fork.

If there are concerns with a resident's nutritional intake or weight, report this to other care staff and document what the resident has eaten and if any difficulties were experienced. Please see our example food chart section on page 27.



Top Tip: If your residents are prescribed an oral nutrition supplement (ONS), **DO NOT** give these at mealtimes. Encourage your residents with their meals and puddings so that they get the maximum nutritional benefit from their meal. ONS should be offered between meals in addition to suitable high calorie snacks and high calorie drinks.

Healthy Eating

Residents who do not have specific nutritional requirements due to illness or disease will require a healthy, balanced diet, in line with the Eatwell Guide (see page 33). For these residents, we aim for a healthy weight which is indicated by a BMI of 20-25kg/m². The Eatwell Guide recommends the following:

Fruit and vegetables – At least 5 portions should be eaten every day including a variety of fruit and vegetables every day. These can be fresh, frozen tinned, dried or small amount of juice (150mls of fruit juice counts as 1 portion).

Starchy carbohydrates – This includes potatoes, bread, rice, pasta and other starchy carbohydrates and should be included at every meal. Choose wholegrain or higher fibre versions with less added fat, salt and sugar.

Protein-rich foods – 2-3 portions should be offered daily. This includes meat, fish, eggs and pulses. Choose lean cuts of meat and minimise the amount of processed meat eaten such as sausages and bacon. It is advised that 2 portions of fish are eaten weekly, 1 of which should be oily like salmon or mackerel. If vegetarian or vegan, residents should be offered a variety of protein containing foods such as beans, lentils, chickpeas, soya products, nuts and nut butters, chia seeds, ground linseed, hemp and pumpkin seeds, vegan meat alternatives. If vegetarian, vegetarian meat-alternatives and free range eggs can be offered.

Calcium-rich foods – 2-3 portions should be offered daily including milk and milk products including cheese and yoghurt. Cheese needs to be animal rennet free if vegetarian. If vegan, offer plant based milks fortified with calcium such as fortified almond, coconut, hazelnut, hemp, oat, rice and soya milks and yoghurts or calcium-set tofu. Foods that contain smaller but notable amounts of calcium include kale, pak choi, okra, spring greens, dried figs, almonds, sesame seeds, sunflower seeds.

Fats and oils - Choose unsaturated oils and use in small amounts. If vegan, choose vegetable oils, fats and spreads.

The government has produced information on providing a healthy diet to care home residents. Nutrients of specific consideration for your residents include:

- Fibre (both soluble and insoluble), to promote regular bowel opening (**note this should only be adopted if the resident is drinking sufficient fluids**).
- Iron, to prevent anaemia. Absorption is enhanced when iron rich foods are consumed alongside foods and drinks rich in vitamin C.
- Calcium, to maintain bone density and prevent osteoporosis development or progression.

Vegetarian and Vegan Diets

Vegetarian and vegan diets are on the rise. The Vegetarian Society (UK) defines a vegetarian as 'someone who lives on a diet of grains, pulses, nuts, seeds, vegetables and fruits with, or without the use of dairy products and eggs. A vegetarian does not eat any meat, poultry, game, fish, shellfish or by-products of slaughter' (www.vegsoc.org). The Vegan Society (UK) defines veganism as 'a way of living which seeks to exclude, as far as is possible and practicable, all forms of exploitation of, and cruelty to, animals for food, clothing or any other purpose'. It is therefore an ethical movement, which has protected status, rather than just being a diet (www.vegansociety.com).

Reasons for following a vegetarian or vegan diet can vary and may include religion, parental or peer influences, animal welfare, environmental issues and health reasons.

As people age, they have a decreased need for energy but increased requirement and/or decreased absorption of some nutrients. This combined with a smaller appetite and possible physical limitations means that a nutrient dense vegetarian or vegan diet should be eaten in line with individual nutritional requirements. There are a number of suitable vegetarian and vegan alternative foods and these are generally found in supermarkets, with more unusual products available in health food shops or online. Appropriate labelling of foods as 'suitable for vegetarians' or 'suitable for vegans' has made it easier to identify appropriate products. If you do not provide in house catering, please check that your caterer provides both vegetarian and vegan diets.

Without careful management, vegans may develop a number of nutritional deficiencies including B12, calcium, fats & omegas, iodine, iron, protein, Vitamin D and zinc. Please ensure that vegan residents have a vitamin and mineral supplement suitable for vegans.

Food Fortification

A fortified diet describes meals, snacks and drinks to which additional nutrients have been added. This is usually achieved by adding foods such as cream, butter, oil, full fat milk, cheese and dried skimmed milk powder. These can be added during food preparation and cooking or added to the finished meal. The aim of food fortification is to enrich foods to provide a higher nutrient density (specifically calories and protein) without increasing the portion size.

Calories are also known as “energy”. Increasing calories can help maintain weight and prevent weight loss..

Protein is fundamental to help with wound healing and maintaining skin integrity, as well as other bodily functions. Protein rich foods that can be used to increase the protein content of meals include dried skimmed milk powder, milk and grated cheese.

A range of vitamins and minerals are essential for health and all bodily functions. If a resident is not eating well and losing weight, it is very likely that they are also missing out important vitamins and minerals. Therefore, in addition to ensuring that the resident is consuming sufficient calories and protein, it is important that they are also encouraged to eat a wide range of foods, including a range of different coloured fruits and vegetables. If the resident chooses to eat a very restrictive diet, they may require an A-Z multivitamin and mineral supplement – available to purchase over the counter.

Residents who are at moderate to high risk of malnutrition (indicated by a MUST score of ≥ 1), or who have a poor appetite, should be offered smaller, fortified meals with nourishing snacks and nourishing drinks between meals (see suggestions on pages 10 and 17). The appetite will not be stimulated when there are long gaps in between eating occasions. A little and often approach is much more successful to increase intake.

Food fortification should be completed on an individual basis to ensure that it is effective or when preparing fortified meals for a number of residents adding the correct amounts of additional ingredients to the batch to maximise fortification per individual portion. For example, adding a few tablespoons of double cream to 10 portions of mashed potatoes would not be adequate food fortification for those 10 residents. Please see some examples of food fortification on page 10. Please note for vegetarian residents, cheese should be animal rennet free.

For more information on food fortification, please watch the following video:

<https://www.youtube.com/watch?v=2tS7fP7aUy8&feature=youtu.be>

Please Note: If a residents' diet is adequately fortified and they are given appropriate higher calorie snacks and drinks throughout the day, **oral nutrition supplements -should only be prescribed for a resident with a MUST score of 2 after a Dietetic assessment.**



Top Tip: Encourage your care staff to sit down in the dining room and eat their meals with your residents. This could encourage residents to eat and drink well.

Food Fortification Ideas

Food (Approximate Quantity)	Approximate Nutrition Provided Before Fortification		Fortified With	Approximate Nutrition Provided After Fortification	
	Calories (kcal)	Protein (g)		Calories (kcal)	Protein (g)
Mashed potato (1 scoop)	45kcal	1g	<ul style="list-style-type: none"> • 1 knob of butter • 1 tablespoon of double cream • 30g grated cheese 	320kcal	8g
Scrambled eggs (2) on hot buttered toast (1 slice)	330kcal	20g	<ul style="list-style-type: none"> • 1 tablespoon of double cream • 30g grated cheese 	525kcal	27g
Custard (small portion) made with whole-milk (100ml)	120kcal	4g	<ul style="list-style-type: none"> • 1 tablespoon of double cream • 1 heaped teaspoon of milk powder 	235kcal	8g
Porridge made with whole-milk (160g)	185kcal	8g	<ul style="list-style-type: none"> • 180ml full fat milk • 1 tablespoon milk powder • 2 tablespoons of double cream • 1 teaspoon of honey 	375kcal	9g
Baked beans or ravioli (150g)	125kcal	7g	<ul style="list-style-type: none"> • 30g grated cheese (match-box size) 	250kcal	15g
Rice/pasta (200g)	250kcal	5g	<ul style="list-style-type: none"> • 2 teaspoons of oil or butter 	300kcal	5g
Serving of vegetables (80g)	15kcal	0g	<ul style="list-style-type: none"> • 1 knob of butter 	90kcal	0g
Fruit canned in syrup (100g)	57kcal	0g	<ul style="list-style-type: none"> • 3 tablespoons of evaporated milk 	125kcal	4g
Cream of Tomato Soup (1/2 400g tin)	200kcal	3g	<ul style="list-style-type: none"> • 1 heaped tablespoon of milk powder • 2 tablespoons of double cream 	400kcal	9g
	<p>Top Tip: Instead of using gravy, look at adding white sauces, cheese sauces, pepper sauces or parsley sauces made with fortified full cream milk (see page 14) to really boost the calorie content of meals for those residents who require a fortified diet.</p>				

Vegan Food Fortification

Items below can be used to fortify foods for example:

50g porridge oats made with 270mls soya milk = **468kcals**

Add 30mls soya cream, ½ sliced banana, 15g peanut butter, 15g syrup = **668kcals**

Please note coconut products are higher in calories but also in saturated fats.

Vegan Alternative		kcal (g/ml)	Protein (g)
Cheese – on sandwiches/toast, stir into baked beans, add to mashed potato, add to soups, on top of jacket potato	Supermarket 'Free From' cheese – 30g Supermarket Free From' soft cheeses – 30g	86 86	0 2
Milk – fortify with soya / oat/coconut cream and make nourishing hot or cold drinks	Alpro Growing Up Milk (soya) – 250mls Oat barista milk – 250mls	163 148	6.3 2.5
Yoghurt – fortify with cream, add to fruit smoothies	Soya fruit yoghurt -125g Soya Greek style yoghurt – 150g Coconut yoghurt – 125g	85 123 164	5 7 2
Ice cream – add to milkshakes, enjoy with fruit or vegan puddings Dessert – fortify with cream, add to milkshakes	Soya – 100g Soya – 125g Coconut – 130g	166 104 150	2.3 4 2
Butter – on toast, sandwiches, tea cakes, fruit loaf add to vegetables and mashed potatoes Use to fry and roast	Vegan spread – 15g Vegetable oil -15ml	80 124	0 0
Cream – add to yoghurts, desserts, smoothies, nourishing drinks, enjoy with fresh fruit	Coconut cream – 30mls Soya cream – 30mls Oat cream – 30g Oat crème fraiche – 30g	66 37 45 53	0 1 0 0
Miscellaneous – on toast, sandwiches, toppings on porridge, pancakes *free from Palm oil	Vegan friendly jam/conserve – 15g Lemon Curd – 15g Nut butter* – 15g Lyle's Golden Syrup – 15g	83 93 90 47	- - 4 0

Nourishing Snacks

Nourishing, nutritious snacks form an important part of a fortified diet. They are a useful source of additional calories and protein and are often considered much more appealing to residents than oral nutrition supplements.

Fruit, low-fat yoghurt and plain biscuits are good snacks for residents with normal dietary requirements. However, these snacks are not sufficient for those residents at moderate to high risk of malnutrition (MUST score ≥ 1) who require additional calories and protein.

To gain weight, residents require approximately at least an additional 600 calories per day from food fortification and nourishing snacks, drinks and puddings in addition to their baseline dietary intake.

Nourishing Snack Ideas

This table outlines nourishing snacks and drinks. For residents who require a high calorie diet, **you should aim for at least 200kcal per snack.**

Snacks suitable for texture modified diets are indicated as follows: soft and bite sized level 6 (previously soft), minced and moist level 5 (previously fork mashable) and puree level 4.

100 calories or less	150 calories
<ul style="list-style-type: none">• Whole-milk 150ml• Full fat yoghurt 150g (5,6,4)• Mini swiss roll x 1• Fully coated chocolate biscuit x 1• Shortbread finger x 1• Digestive biscuit x 1• Rich Tea biscuit x 2	<ul style="list-style-type: none">• Thick and creamy yoghurt 150g (5,6,4)• Crumpet x 1 with butter• Scotch pancake x 1 with butter• Ready-to-eat custard pot 150g (5,6,4)• Ready-to-eat rice pudding 150g (5,6)• Chocolate coated ice cream block (5,6)
200 calories	250 calories
<ul style="list-style-type: none">• Fruit corner type yogurt (5,6)• Individual ready-to-eat trifle (5,6)• Dairy ice-cream x 2 scoops (5,6)• Cereal (30g) with whole-milk (100g) (5,6)• Cream cracker x 2 with butter and cheddar cheese (30g)• Digestive biscuits x 2 with butter• Oatcakes x 2 with butter• Crumpet x 1 with butter and jam• Scotch pancake x 1 with butter and jam• Fruit malt loaf slice x 1 with butter	<ul style="list-style-type: none">• Small scone x 1 with butter• Oatcakes x 2 with butter and cheese• Ring doughnut x 1• Jam doughnut x 1• Ready-to-eat rice pudding 200g (5,6)
300 calories	
	<ul style="list-style-type: none">• Scone x 1 with butter and jam• Chocolate 50g• Digestive biscuits x 2, butter and cheese• Enriched whole-milk (half a pint with 2 tbsp. dried skimmed milk powder)

Nourishing Drinks

Nourishing drinks are a tasty way to increase the calories for residents who are at risk of malnutrition. It is important to encourage your residents who are at risk of malnutrition have at least 2 nourishing drinks per day instead of their usual cup of tea or cup of coffee – standard cups of tea and coffee have very few calories. Here are few tried and tested recipes that are much more beneficial than their unfortified versions. **Most of these nourishing drink recipes have more calories than ONS.**

Fortified Milk – NB NOT APPROPRIATE FOR CKD STAGE 4 & 5

Ingredients

- 1 pint of full cream milk
- 4 heaped tablespoons of skimmed milk powder (approx. 65g)

Method

Mix a splash of full fat milk with the milk powder to form a paste with a fork. Whisk the rest of milk into the paste and continue to whisk with a fork until smooth. Use in any cereals, porridge, hot and cold drinks etc.



Approx 600kcal + 44g protein

Hot Chocolate

Ingredients

- 150ml full fat milk
- 4 teaspoons (16g) of skimmed milk powder
- 4 tablespoons (60ml) of double cream
- 4 teaspoons (15g) of hot chocolate powder

Method

Mix all of the ingredients together and heat to the desired temperature. Add extra sugar if required.



Approx: 500kcal + 13g protein

Milkshake

Ingredients

- 200ml full fat milk
- 2 heaped tablespoons (30g) of skimmed milk powder
- 1 tablespoon (15ml) of double cream
- 4 teaspoons (20g) milkshake powder with added vitamins and minerals e.g. Nesquik, Asda, Lidl or Morrison's milkshake mix

Method

Whisk all of the ingredients together. Serve chilled.



Approx: 315kcal + 18g protein

Milky Coffee

Ingredients

- 200ml full fat milk
- 1 heaped tablespoon (15g) of skimmed milk powder
- 1 teaspoon (5g) of sugar
- 1 teaspoon (2g) of coffee

Method

Mix all of the ingredients together and heat to the desired temperature.
Add extra sugar if required.



Approx: 200kcal + 12g protein

Fortified Cup-A-Soup

Ingredients

- 1 cup-a-soup sachet
- 200ml full fat milk
- 1.5 tablespoons (20g) of skimmed milk powder



Method

Whilst warming the milk to the desired temperature, add the milk powder into the milk and stir well.
Gradually add the cup-a-soup sachet to the warm milk and again stir well.

Approx: 280kcal + 15g protein

Peaches & Cream Smoothie

Ingredients

- 100g full-fat Greek yoghurt
- 100g of tinned peaches in syrup
- 1 heaped tablespoon (15g) of skimmed milk powder
- 100ml full fat milk
- 1 tablespoon (15ml) double cream

Method

Mix all the ingredients together in a blender blitz until smooth – ensure there are no lumps.

Approx: 370kcal + 14g protein



Dairy Free Chocolate, Banana & Peanut Butter Smoothie

Ingredients

- 180ml sweetened soya milk
- 1 heaped tablespoon (25g) smooth peanut butter
- 1 teaspoon (5g) cocoa powder
- 1 teaspoon (4g) vegetable oil
- 4 teaspoons (20g) sugar / honey
- Half a ripe banana (50g)

Method

Mix all the ingredients together in a blender blitz until smooth – ensure there are no lumps.



Approx: 420kcal + 14g protein

Fortified Malty Drink

Ingredients

- 200ml full fat milk
- 2 heaped tablespoons (30g) skimmed milk powder
- 5 heaped teaspoons (25g) Ovaltine Original / Horlicks drink

Method

Mix all of the ingredients together and heat to the desired temperature.
Add extra sugar if required.

Approx: 330kcal + 20g protein

Fruit Fizz

Ingredients

- 100ml fruit juice
- 100ml lemonade (not 'sugar free' or 'no-added sugar')
- 30ml high-juice squash (not 'no-added sugar')
- 1 tablespoon (15g) sugar
- 1 scoop (50g) vanilla ice-cream

Method

Add the fruit juice, high-juice and lemonade into a tall glass and mix well.

Add the sugar and mix well.

Add the ice-cream and mix well.

Approx: 250kcal + 2g protein



Top Tip: Add grated cheese to the top of Sheppard's Pies, Cottage Pies and pasta bakes before baking.

Nourishing Puddings

Nourishing puddings are an excellent way of increasing a residents' calorie intake and can be used as snacks or after main meals. As well as fortifying custard or using high calorie accompaniments such as cream or ice-cream to go alongside puddings, small high calorie puddings can be made easily for residents requiring a fortified diet.

Sweet Milk Jelly – Makes 4 servings

Ingredients

- 1 x 135g block of jelly (**NOT** no-added sugar or sugar-free)
- ½ pint (285ml) boiling water
- ½ pint (285ml) sweetened condensed milk

Method

Split jelly into cubes and place in a jug / bowl. Add the boiling water and stir until dissolved. Add the condensed milk and stir; then pour into moulds / serving dishes. Allow to cool and then refrigerate until set.

Per portion approx: 320kcal + 7g protein

Fortified Instant Whip – Makes 3 servings

Ingredients

- 200ml full fat milk
- 100ml double cream
- 3 heaped tablespoons (40g) skimmed milk powder
- 1 packet of instant whip (**NOT** sugar free). Either Angel Delight or supermarket brand

Method

Whisk together the milk powder and milk. Whisk in the double cream. Whisk in the instant whip – whisk well. Split into 3 serving dishes and leave to thicken.

Per portion approx: 320kcal + 8g protein

Citrus Cream – Makes 4 servings

Ingredients

- 300ml double cream
- 70g caster sugar
- Zest and juice of 1 orange or lemon
- 2 heaped tablespoons (30g) skimmed milk powder

Method

Put cream and milk powder in a saucepan and heat until the milk powder has dissolved. Stir in the sugar and simmer whilst stirring for a minute. Take off the heat and stir in the zest and juice. Pour into 4 serving dishes and chill.

Per portion approx: 450kcal + 4g protein



Top Tip: If you have a resident who is struggling with, has gone off or is refusing savoury foods, additional portions of puddings such as the ones above can be given to increase their calorie and protein intake.

Example Fortified Menu

Food fortification can be a powerful tool to increase the calorie and protein content of a malnourished resident's diet. Below are two menus (based on an average older residents' intake) - one with and one with fortification. The difference between the days menu and be seen and compared.

	Unfortified Menu	Fortified Menu
Breakfast	Porridge made with full fat milk and a cup of tea	Fortified Porridge with a cup of tea made with fortified milk
Mid - Morning	2 rich tea biscuits and a coffee with full fat milk	Fruit malt loaf slice x 1 with butter and a fortified milky coffee
Lunch	Small portion of Shepherd's Pie with mixed vegetables and glass of squash	Small portion of fortified Shepherd's Pie and fortified mixed vegetables with a glass of squash
Mid – Afternoon	1 shortbread finger and a cup of tea	1 scone with butter and jam and a cup of tea
Evening Meal	Baked beans on 1 slice of toast and Instant Whip and a glass of squash	Cheese and baked beans on a slice of buttered toast and a fortified Instant Whip and a glass of squash
Super	Malted drink	Fortified malty drink

Calories + Protein	1,100kcal + 49g protein	2,580kcal + 97g protein
---------------------------	--------------------------------	--------------------------------

The fortified menu contains 1480kcal and 48g protein more than the non-fortified menu.

With 2 compact ONS the non-fortified menu would only increase to 1700kcal and 73g protein – **the fully fortified menu without ONS still 880kcal and 24g protein more than the non-fortified menu with oral nutrition supplements.**

Food fortification will require catering, carers, nurses and management staff to work together to ensure that residents receive their individualised nutrition and hydration needs in line with CQC regulation 14.

	Top Tip: Protein is incredibly important for your residents therefore it is important that protein is distributed throughout the day to enable them to achieve their protein requirements. Traditional breakfast options such as toast or cereal do not help residents to do this. Breakfasts such include options such as baked beans, eggs and other cooked breakfast items. Ideas can be found at: https://www.malnutritionpathway.co.uk/library/proteinideas.pdf .
-------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Eating Well With Dementia

Individuals with dementia may experience reduced or limited recognition of hunger and / or thirst. Consequently, this can create anxieties and can bring with it challenges at mealtimes. It is important to try and make mealtimes as relaxed as possible. As the environment is important when eating and drinking it should ideally be free from noise and distractions. Individuals may have difficulty recognizing foods and with eating and drinking.

Here are tips to help individuals with Dementia:

- Turn off any nearby televisions or radios.
- Use plain plates which are bright in colour (e.g. red / blue / yellow) as it provides a greater contrast between the food and the plate.
- Use a plain coloured tablecloth so cutlery and food stand out. Avoid bright patterns as these can be confusing.
- Avoid distractions on the table such as plants, ornaments, books, magazines and condiments.

For Residents Unable To Sit Down For A Meal

- Offer finger foods, in line with any Speech and Language Therapy recommendations, such as, sandwiches, chips, croquettes, fish fingers, sausage rolls, pizza, quiche, sliced vegetables, cheese and crackers, scones, cakes or pieces of fruit.
- Have readily available snacks for individuals to have whilst wandering / on the go.
- Join individuals at the table and provide encouragement and the opportunity for them to 'model' your behavior.
- Consider changing mealtimes where people are more likely to eat at different times.
- Offer food and drinks and fluids using the 'little and often' approach.

For Residents Not Able To Recognise / Use Cutlery

- Try helping the individual by placing cutlery in their hands and guiding it to their mouth hand over hand.
- Try eating with the individual so they can watch as this can be a helpful prompt or reminder.
- Don't worry about individuals eating foods with their fingers if they are struggling to use cutlery.

For Residents Spilling Food / Drink

- If an individual has difficulty using a knife, try chopping up food for them.
- Prepare foods that are easy to eat with a spoon.
- Try not to overfill cups and if necessary offer drinks more frequently.
- Use place mats and table cloths that can be wiped clean.
- An occupational therapist may be able to provide advice about adapted crockery and cutlery.

For Residents With Changes To Food Preferences

- Don't worry if individuals seem to have altered food preferences or unusual food combinations as they may be unsure of what food items typically go together.
- Support them to eat the foods they like even though they may have disliked them

previously.

For Residents Not Finishing Meals

- Give the individual plenty of time to eat. Verbal prompts such as, 'How is your meal? / Would you like a drink?' may provide encouragement.
- Encourage 6 small meals throughout the day rather than 3 main meals and include nourishing snacks and drinks.
- Fortifying foods with for example cheese, butter or sugar to increase their nutrient content is outlined in a leaflet called Big Nutrition which is available from your local Dietitians or GP.
- Be aware that the consistency of food / drink an individual is able to manage may change due to problems with chewing / swallowing. They may start to hold food in their mouth, chew more than before or refuse to eat foods that are harder to chew (e.g. hard vegetables). A doctor or speech and language therapist may be able to provide support and advice about this.
- If the individual is eating less and is losing weight without trying speak to their GP or get advice from a dietitian.

For Residents Not Drinking Enough

Sometimes the person with dementia may not recognise that they are thirsty or may forget to drink.

- Offer drinks frequently.
- Include drinks in social interactions e.g. having a cup of tea and a chat.
- Some individuals may need support to hold a cup / place the cup gently in their hand. Tell them what drink is in the cup.
- Try using clear cups so that individuals can see what they are drinking and if they have a favorite cup, use this.
- Provide verbal prompts such as, 'how's your tea' or simple written / picture notes about drink e.g. a picture of milk on the fridge. Leaving drinks / jugs of fluids out can also act as a visual prompt.
- Encourage a variety of fluids including both warm and cold drinks.
- An occupational therapist maybe able to help with non-spill cups if spilling is a problem.

Supporting Dementia Residents To Eat And Drink

If the person you care for requires support at and drink:

- Sit at eye level or slightly below the individual. You may find it helpful to sit at one side or slightly in front of them and maintain eye contact.
- Talk about the food and drink that you are offering, but try to discourage the person from talking whilst eating because of the risk of choking.
- Naming foods and drinks can help trigger memories.

Diabetes

Type 1 diabetes is a lifelong condition and is often diagnosed in childhood. The body attacks the cells in the pancreas that make insulin so the body then becomes unable to produce its own insulin. The exact cause of this is not yet known. Our bodies need insulin as it allows glucose to enter our cells to be used for energy. People with Type 1 diabetes will require insulin injections. Their insulin regimen will be advised by their diabetes consultant / specialist nurse to meet their requirements.

Type 2 diabetes is often diagnosed in later life. The pancreas releases insulin in response to a rise in blood glucose however the insulin does not work effectively. The blood glucose level continues to rise and the pancreas tries to release more insulin in response. Eventually some people with Type 2 diabetes can tire their pancreas out, resulting in less and less insulin being produced. People with Type 2 diabetes may be diet controlled or have medications prescribed which could include Metformin, Gliptins, Gliflozins, insulin injections as well as others.

All individuals with Type 1 and 2 diabetes will have their long-term blood glucose management monitored by their GP surgeries. Here are 10 tips for eating well with diabetes from Diabetes UK. **Please note: if a resident needs to gain weight with diabetes, they should have a fortified diet, snacks and high calorie drinks as per pages 10 to 17. Their diabetes team may need to alter their medications to accommodate their diet if malnourished. The sugar can be reduced in the fortified recipes if required (this will however lower the calorie content).**

1. Choose healthier carbohydrates

All carbohydrates affect blood glucose levels. It is important to choose healthy carbohydrate sources and cut down on foods low in fibre such as white bread, white rice and white pasta – brown and wholegrain varieties are better alternatives. Carbohydrate portion sizes also require consideration.

2. Eat less salt

Eating lots of salt can increase the risk of heart disease and stroke. Those with diabetes are already more at risk of these conditions. Aim for a maximum of 6g (one teaspoonful) of salt a day in your residents' diet. A lot of pre-packaged foods already contain salt so remember to read food labels and choose those with less salt. Cooking from scratch will help to reduce the amount of salt consumed. Get creative try to reduce your residents' salt intake by trying different herbs and spices to add flavour.

3. Eat less red and processed meat

It is best to avoid foods such as ham, bacon, sausages, beef and lamb on a regular basis. These have been linked to heart problems and cancers. Try swapping red and processed meat in care home menus for pulses such as beans and lentils or eggs, fish, poultry e.g. chicken and turkey or unsalted nuts. Beans, peas and lentils are very high in fibre and don't affect blood glucose levels significantly. Try and include two portions of oily fish a week e.g. mackerel or salmon.

4. Eat more fruit and veg

It is important for residents to eat a variety of fruits and vegetables – 5 portions or more daily. Fruits contain natural sugar and should not be avoided. This is different to added sugar (also known as free sugars) found in foods such as chocolate, biscuits and cakes. Products like fruit juices also count as free sugar, so go for whole fruit instead. Fruit can be fresh, frozen, dried or tinned (in juice, not in syrup) - it all counts towards the 5 a day. Fruit is best eaten throughout the day instead of larger portions.

5. Choose healthier fats

Fats are needed in the diet but different fats affect our health in different ways. Healthier fats are found in foods such as unsalted nuts, seeds, avocados, oily fish, olive oil, rapeseed oil and sunflower oil. Some fats can increase the amount of cholesterol in your blood, increasing the risk of heart problems. These are mainly found in animal products and prepared foods, e.g. red and processed meat, ghee, butter, lard, biscuits, cakes, pies and pastries. It is a good idea to cut down on using oils in general, so try to grill, steam or bake foods instead of frying. For those residents needing to gain weight with diabetes, food should still be fortified – see our food fortifications sections.

6. Cut down on added sugar

Swapping sugary drinks, energy drinks and fruit juices should be switched for water, milk, tea or coffee without sugar, and low calorie soft drinks. Cutting out added sugars can help control blood glucose levels and help weight management if a person is overweight. Low or zero calorie sweeteners (also known as artificial sweeteners) can be used in drinks to add sweetness without sugar.

7. Be smart with snacks

For those wanting to maintain a healthy weight or lose weight, choose yoghurts, unsalted nuts, seeds, fruits and vegetables instead of crisps, chips, biscuits and chocolates. For those needing to gain weight, see our suggested snacks list to increase calories.

8. Drink alcohol sensibly

Alcohol is high in calories. Try to keep residents' to a maximum of 14 units a week. Taking alcohol on an empty stomach can make hypos more likely to happen if treated Gliclazide or insulin.

9. Don't bother with so-called diabetic food

To say food is a 'diabetic food' is now against the law. This is because there isn't any evidence that these foods offer a special benefit over eating healthy. They can also often contain just as much fat and calories as similar products, and can still affect blood glucose level. These foods can also sometimes have a laxative effect.

10. Get minerals and vitamins from foods

There is no evidence that mineral and vitamin supplements help you manage diabetes. It is advised that supplements are not taken unless advised by a health care professional. Vitamin D should still be taken – see our Vitamin D section. It is better to get your essential

nutrients by eating a mixture of different foods. This is because some supplements can affect your medications or make some diabetes complications worse, like kidney disease.

Don't forget to keep moving

Being more active goes hand in hand with eating healthier. It can help you manage your diabetes and also reduce the risk of heart problems. This is because it increases the amount of glucose used by muscles and helps the body use insulin more efficiently. 150 minutes of moderate intensity activity a week is advised but this may not be possible for some residents. Speak with your GP or local physiotherapy team to see what exercises would be suitable for your care home residents of varying abilities.

Hilos

If a resident experiences a hypo, you can use sugary drinks to treat them followed by a starchy snack to sustain their blood glucose. If a resident is experiencing regular hypos it is important to discuss this with their diabetes team.

Vitamin D

Vitamin D is also known as the ‘sunshine vitamin’. Vitamin D helps to absorb and regulate the amount of calcium and phosphate in the body and is needed to keep bones, teeth and muscles healthy. Our bodies create the active form of vitamin D from exposure to the sun. From late March to the end of September, between 11am-3pm, the sunlight (specifically UVB rays) is strong enough for people to synthesise Vitamin D from having their forearms, hands and lower legs uncovered for short periods. Care should be taken not to burn in the sun; therefore sunscreen should be worn before the skin starts to turn red or burn.

There are several factors that affect the amount of active vitamin D produced in the skin. Those with darker skin colours will need to spend longer in the sun to produce the same amount of vitamin D than someone with lighter coloured skin.

Vitamin D can also be found in a small number of foods including:

- oily fish such as salmon, sardines, herring and mackerel
- red meat
- liver
- egg yolks
- fortified foods such as fats, spreads and some breakfast cereals.

In the UK, cow’s milk is not considered a good source of vitamin D because it is not fortified as it is in some other countries. It is unlikely that our vitamin D requirements can be met via diet alone.

Residents in care homes often do not get adequate exposure to the sun for vitamin D synthesis. Sitting by a window is not adequate for vitamin D synthesis because the glazing blocks the UVB rays required. Many care home residents may therefore be deficient in vitamin D. If resident’s become deficient in Vitamin D, they may experience the following:

- Chronic fatigue and tiredness
- Back, hip and bone pain
- Depression
- Impaired wound healing
- Hair loss
- Muscle pain

It is advisable that care home residents take a Vitamin D supplement of 10 micrograms / 400IU daily. These can be purchased from supermarkets or local pharmacies by care homes or residents’ families. If long term deficiency is suspected, a blood test may be required via the residents’ GP and prescribed high dose Vitamin D may be required for a short period. It is then advised that residents take a maintenance dose of Vitamin D (20 micrograms / 800IU daily) and these can be purchased from supermarkets or local pharmacies by care homes or residents’ families.

For residents following a vegan diet, vitamin D3 supplements from lichen sources (vegan friendly) are available from various online retailers or health food shops – again these can be purchased by care homes or residents’ families.

Coeliac Disease

Coeliac disease is a serious illness where the body's immune system attacks its' own tissues when gluten is eaten. This causes damage to the gut lining, meaning that the body can't properly absorb nutrients from food. Coeliac disease is not an allergy or food intolerance. Coeliac disease is common and affects around one in 100 people. Only 30% who have the condition have been diagnosed meaning there are currently around half a million people in the UK who have coeliac disease but aren't aware. If a first degree family member (such as mother, father, sister or brother) has the condition then the chances of having it increase to one in ten. Dermatitis herpetiformis (DH) is the skin manifestation of coeliac disease which occurs as a rash that commonly occurs on the elbows, knees, shoulders, buttocks and face, with red, raised patches often with blisters.

Symptoms range from mild to severe and can include bloating, diarrhoea, nausea, wind, constipation, tiredness, mouth ulcers, sudden or unexpected weight loss (not in all cases), and anaemia. Once diagnosed, the only treatment for coeliac disease is a gluten free diet. Gluten is found in wheat, barley and rye. Some people are also sensitive to oats. 100% compliance to a gluten free diet is advised to avoid later complications with Coeliac Disease.

Naturally gluten free foods include:

- Meat
- Fish
- Fruit and vegetables
- Rice
- Potatoes
- Lentils

Breads and bakery products, cereals etc. need to be gluten free versions – these are no longer available on prescription in Derbyshire and have to be purchased privately either at supermarkets or online.

Catering for individuals with Coeliac Disease requires careful consideration to avoid cross contamination. Here are tips to avoid cross contamination:

- Wipe down surfaces and ideally prepare gluten free foods first.
- Clean pots and pans with soap and hot water. Washing up liquids are fine to use and standard washing up liquid or using a dishwasher will remove gluten. You do not need to use separate cloths or sponges.
- Have separate bread boards to keep gluten free and gluten containing breads separate.
- Use a separate toaster or toaster bags – breadcrumbs in toasters are a source of contamination.
- Use clean oil or a separate fryer for frying gluten free foods.
- Have different butter / jams etc. to prevent contamination or use spoons and a 1 dip rule to prevent breadcrumbs from getting into condiments.

Hydration

Drinking adequate fluid is fundamental to preventing dehydration and can also help to prevent urinary tract infections, acute kidney injuries, constipation, pressure ulcers, poor wound healing and cognitive impairment. In addition, adequate hydration is beneficial in the management of heart disease by protecting against clot formation, can help to maintain healthy blood glucose levels and is an important part of any multifactorial falls prevention strategy.

The UK Government recommends that 6-8 glasses of water (at least 1 to 2 litres), or other fluids, should be consumed every day.

The following tips can help you to promote adequate hydration in your residents:

- Ensure that all residents have a jug of fresh water and a glass within their reach at their bedside.
- Encourage staff to prompt all residents with their fluid intake throughout the day; some residents may not recognise the feeling of thirst.
- Ask family members and others to prompt the resident to drink whilst they visit.
- Offer a drink at every meal; take the time to find out what a resident likes to drink by asking them or their family.
- Residents often drink all of the fluid in their glass when taking medications; therefore offer a larger volume at this occasion to encourage them to drink more.
- On warmer days or in warm rooms we all need to drink more than normal.
- Residents with dementia can often see fluids better when they are in a dark cup. Research has found this has increased fluid intake for individuals with this medical condition.
- For residents who require thickened fluids, ensure their drinks are all thickened to the correct descriptor.

Food Charts

Food charts can be a very important tool for you and Dietitians to look at how much a resident is eating and drinking. It helps to identify trends, and if completed accurately, can help a dietitian to calculate how many calories and grams of protein your resident is consuming. These do not need to be kept for all residents. We recommend a 3 day food chart for residents who you are concerned about or in the run up to a Dietitian's assessment. Here is an example of the details which should be recorded.

Resident's Name: Mr Bob Smith

Date: 1st May 2021

Meal	Food Offered	Portion Size	Amount Eaten
Breakfast	Porridge - fortified with 1 tspn of honey, 1 tbsp of double cream and 2 tspns of milk powder + Milky coffee (200ml fortified milk)	Medium Bowl	3/4 + Approx 125ml
Mid - Morning	2 x chocolate digestives + cup of tea 200ml (made with fortified milk) + Fortisip Compact 125ml	-	All + 1/2
Lunch	Cottage Pie Carrots and Bean with a teaspoon of butter added + Swiss Roll and double cream (approx. 100ml) + Orange Squash 200ml	Small - tea plate portion + Approx inch slice	All + All + 1/2

Over The Counter Oral Nutrition Supplements

Some ONS, such as Complan, Complan Smoothie, Complan Soup or Meritene Shake, Meritene Soup and Meritene Ready To Drink, can be purchased over the counter (OTC). These might be appropriate for residents who remain at high risk of malnutrition (MUST of 2 or greater) despite being offered a well-fortified diet as per their nutritional care plan.

Oral Nutrition Supplements Which Can Be Prescribed

The Derbyshire Management of undernutrition in adults Guidelines recommend that individuals who remain at high risk of malnutrition, despite being offered a well-fortified meals, high calorie snacks, drinks and puddings and have used OTC ONS for four weeks, may need to trial ONS.

Important points to remember about ONS:

- They are intended for short-term use for residents with a high malnutrition risk (≥ 2).
- They should not replace fortified meals, high calorie snacks or high calorie drinks. They should be given **in addition** to a fortified diet and offered in between mealtimes.
- Once opened, ONS can be kept safely for up to 4 hours at room temperature or for up to 24 hours in the refrigerator. After this time has elapsed, dispose of the opened ONS.
- ONS are prescribed; therefore the amount consumed **must** be recorded.
- ONS should only be given to the intended resident, as they could cause adverse effects in another resident.
- The prescription of ONS should be advised by a Dietitian and residents must meet the ACBS Criteria **and** local prescribing guidelines to qualify for an NHS prescription.
- ONS **should not** be prescribed instead of a fortified diet and high calorie snacks.

The Derbyshire Oral Nutrition Support Guidelines for Adults can be seen at: [BNF 9 Nutrition & Blood](#)

Referral to Nutrition and Dietetics

Your local Community Nutrition and Dietetic Services are based at the Chesterfield Royal Hospital (who cover North Derbyshire) and Florence Nightingale Community Hospital, Derby (who cover South Derbyshire). Referrals are accepted for a range of different conditions.

If you wish to discuss whether a referral is appropriate or wish to request a referral form, do not hesitate to contact the departments at crhft.nutritionanddietetics@nhs.net (Chesterfield Royal Hospital) or dhft.cnd@nhs.net (Florence Nightingale Community Hospital, Derby) or on the telephone numbers on the front cover of this pack.

When referring a resident for Nutrition Support advice, please provide the following information as a minimum and any other information which you think might be helpful:

- Current weight (kg)
- Height (m)
- Weight history from the last 6 months
- Current MUST score
- Reason for referral
- Diagnosis or relevant past medical history
- Relevant medications
- Actions already taken to minimise weight loss or improve dietary intake

Resource Section



Monthly Catering Communication Sheet

Good communication is essential between all staff members to ensure that residents receive the correct meals for their nutritional needs. Consideration needs to be given to textures, calorie and protein content and other dietary issues such as allergies. This form can record the dietary needs of each of each of your residents, to support your catering staff in providing personalised meals to meet your residents' requirements.

Residents Name		MUST Score (0 = low, 1 = med, 2+ = high)	Food Texture	Dietary Requirements (e.g. nutrition support, weight reduction)	Allergies (Yes / No) If yes, please state
1		High / Med / Low			
2		High / Med / Low			
3		High / Med / Low			
4		High / Med / Low			
5		High / Med / Low			
6		High / Med / Low			
7		High / Med / Low			
8		High / Med / Low			
9		High / Med / Low			
10		High / Med / Low			
11		High / Med / Low			
12		High / Med / Low			
13		High / Med / Low			
14		High / Med / Low			

15		High / Med / Low			
16		High / Med / Low			
17		High / Med / Low			
18		High / Med / Low			
19		High / Med / Low			
20		High / Med / Low			
21		High / Med / Low			
22		High / Med / Low			
23		High / Med / Low			
24		High / Med / Low			
25		High / Med / Low			
26		High / Med / Low			
27		High / Med / Low			
28		High / Med / Low			
29		High / Med / Low			
30		High / Med / Low			

Month:	Date Completed:
Completed By:	Job Title:

Eatwell Guide

Check the label on packaged foods

Each serving (150g) contains			
Energy 104kJ 25kcal	Fat 3.0g	Sugars 1.3g	Salt 0.8g
13%	4%	7%	15%

of an adult's reference intake
Typical values (as sold) per 100g: 887kJ / 210kcal

Choose foods lower in fat, salt and sugars

Use the Eatwell Guide to help you get a balance of healthier and more sustainable food.
It shows how much of what you eat overall should come from each food group.

Eat at least 5 portions of a variety of fruit and vegetables every day



Eat at least 5 portions of a variety of fruit and vegetables every day



Eat more beans and pulses, 2 portions of sustainably sourced fish per week, one of which is oily. Eat less red and processed meat



Eat less often and in small amounts



Choose wholegrain or higher fibre versions with less added fat, salt and sugar

Potatoes, bread, rice, pasta and other starchy carbohydrates



Water, lower fat milk, sugar-free drinks including tea and coffee all count.
Limit fruit juice and/or smoothies to a total of 150ml a day.



Oil & spreads
Choose unsaturated oils and use in small amounts

Per day 2000kcal
2500kcal = ALL FOOD + ALL DRINKS

Source: Public Health England in association with the Welsh Government, Food Standards Scotland and the Food Standards Agency in Northern Ireland

© Crown copyright 2016

Algorithm for Nutrition Care Planning

ALGORITHM FOR NUTRITION CARE PLANNING

The Malnutrition Universal Screening Tool (MUST) score should be completed for all* individuals on admission to hospital (and as minimum weekly) or at first contact (and as a minimum monthly if in community). The details of this should be documented on MUST.

*If an individual is approaching the end of their life, MUST scores will no longer be appropriate however you may wish to provide some elements of this care plan.

The action taken for each of these will differ depending upon if the healthcare professional is supporting the individual with eating and drinking (as for example when they are in hospital) or if they are caring or supporting an individual who is living at home and may have support with eating and drinking from carers.

All of the following should be considered and included as required in the nutrition care plan for all individuals irrespective of MUST score.	
MUST score 0	<ul style="list-style-type: none">Provide / support individual / carers to provide a diet that meets all ethnic and other dietary requirements (e.g. gluten free diet, diet for diabetes, texture modified diet).Support / encourage carers to allow individuals to make their own food choices showing them foods or pictures if required.
MUST score 1	<ul style="list-style-type: none">Offer alternatives if individuals refuse a food / meal, encourage a 'little and often' approach and / or offering finger foods.Aim not to create a stressful environment for individuals if they refuse a meal or are unable to eat.Provide / support individuals to access all necessary aids to maximise eating and drinking (e.g. utensils, dentures, glasses, hearing aids, adapted feeding utensils¹).Assess / ask about dentition² and oral hygiene and provide / promote good oral hygiene.Provide assistance / advise carers on how to provide assistance to individuals with food choices and feeding.
MUST score 2	<ul style="list-style-type: none">Address any underlying medical³ (e.g. gastrointestinal symptoms, pain), physical (e.g. dysphagia⁴, unable to put food to mouth, problems with chewing) social⁵ (e.g. isolation, access to food, finances,) or emotional (e.g. bereavement, mental health issues⁶, depression) causes of poor appetite / food refusal.Promote appropriate position⁷ for eating and drinking, encourage social eating and a pleasant homely atmosphere for eating.Offer / recommend hand washing or wipes for personal hygiene before meals.Ensure meals presented attractively, at the correct temperature, and within reach where appropriate.
MUST score 1 Medium risk Observe	<p>Continue with all of the above and consider and include the following as required.</p> <ul style="list-style-type: none">Agree realistic aims and outcomes with the individual / carers (e.g. increased nutritional intake).Encourage and offer nutritious snacks and milky drinks in between mealsConsider introducing / promoting over the counter oral nutrition supplement e.g. Meritene Energis, Complan between meals (twice daily maximum) for those without CKD 4 / 5.Provide fortified meals or give written information to enable individual to fortify their own meals (e.g. Big Nutrition for Small Appetites).Keep / support individuals or carers to keep a 3 day food and drink record chart.Review and monitor compliance and progress towards the agreed aims and outcomes.Review aims and outcomes and adapt nutrition care plan as required.
MUST score 2 High risk Treat	<p>Continue with all of the above and consider and include the following as required.</p> <ul style="list-style-type: none">Agree realistic aims and outcomes with the individual / carers (e.g. increased nutritional intake).Consider one month trial of prescribable oral nutritional supplements.Consider a referral to the dietitian.

Referrals to other services may be required as indicated by the footnotes.

¹ Occupational Therapist

² Dentist / Oral Health Team

³ Medical team or Advanced Nurse / Clinical Practitioner

⁴ Speech and Language Therapist


⁵ Social services

⁶ Mental health team

⁷ Physiotherapist

Texture Modified Diet, Fluids and IDDSI

The International Dysphagia Diet Standardisation Initiative (IDDSI) framework consists of a continuum of 8 levels (0 - 7), where drinks are measured from Levels 0 – 4, while foods are measured from Levels 3 – 7. The IDDSI Framework provides a common terminology to describe food textures and drink thickness.



It is important to use the scoop provided with the thickener.

IDDSI Level of Thickness	Number of Scoops for Nutilis Clear	Number of Scoops for Thick & Easy Clear
1	1	1
2	2	2
3	3	3
4	7	6

The Speech and Language Department at Chesterfield Royal have put together a helpful video to demonstrate how the IDDSI descriptors should be mixed and how to make them more enjoyable. It can be found at: <https://www.youtube.com/watch?v=hzdXykWPrXE>

or search YouTube for ' Chesterfield Royal Hospital How to Thicken Fluids'



Food also has IDDSI descriptors which can be seen below:

Descriptor	Description	Testing Thickness
0 - Thin	<ul style="list-style-type: none"> Flows like water Fast flow Can drink through any straw, spout, as is appropriate for the person 	Test liquid flows through a 10ml slip tip syringe completely within 10 seconds, leaving no residue
1 – Slightly Thick	<ul style="list-style-type: none"> Thicker than water Requires a little more effort to drink than thin liquids Flows through a straw 	Test liquid flows through a 10ml slip tip syringe leaving 1-4ml in the syringe after 10 seconds
2 – Mildly Thick	<ul style="list-style-type: none"> Flows off a spoon Sippable, pours quickly from a spoon, but slower than thin drinks Effort required to drink through a straw 	Test liquid flows through a 10ml slip tip syringe leaving 4-8ml in the syringe after 10 seconds
3 – Liquidised / Moderately Thick	<ul style="list-style-type: none"> Can be drunk from a cup Some effort is required to suck through a straw Can't be piped, layered or moulded on a plate Can't be eaten with a fork as will drip slowly through the prongs in dollops Can be eaten with a spoon No oral processing or chewing required Smooth texture with no bits (lumps, fibres, husk, gristle etc.) 	Test liquid flows through a 10ml slip tip syringe leaving >8ml in the syringe after 10 seconds Or Drips through prongs of a fork in dollops and spreads out on a flat surface
4 – Pureed / Extremely Thick	<ul style="list-style-type: none"> Usually eaten with a spoon but a fork is possible Can't be drunk from a cup Can't be sucked through a straw 	The prongs of a fork can make a clear pattern in the surface which is retained

	<ul style="list-style-type: none"> • Does not require chewing • Can be piped, layered or moulded • Slow movement under gravity but can't be poured • Falls off spoon in a single spoonful and holds shape • No lumps • Not sticky • Liquid must not separate from the solid 	<p>No lumps</p> <p style="text-align: right;">+</p>
5 – Minced & Moist	<ul style="list-style-type: none"> • Can be eaten with a fork or spoon • Could be eaten with chopsticks in some cases • Can be scooped and shaped • Soft and moist with no separate liquids • Lump size 4mm • Lumps are easy to squash with the tongue 	<p>When pressed with a fork, the particles should easily be separated and come through the prongs</p> <p style="text-align: right;">+</p> <p>Can be easily mashed with little pressure from the fork</p>
6 – Soft & Bite Sized	<ul style="list-style-type: none"> • Can be eaten be a spoon or fork • Can be mashed / broken down with pressure from a fork or- spoon • A knife is not required to cut this food • Chewing is required before swallowing • Soft, tender and moist throughout, but no separate thin liquid • Bite sized pieces for adults 15mm / 1.5cm 	<p>Pressure from a fork held on its side can be used to cut or break this texture into smaller pieces. If one of the lumps are pressed with enough force to turn a thumb nail white, the lump changes shape and doesn't return to its original shape.</p>
7 – Easy to Chew	<ul style="list-style-type: none"> • Normal, everyday foods of soft/tender textures that are developmentally and age appropriate • Any method may be used to eat these foods • Does not include: hard, tough, chewy, fibrous, stringy, crunchy, or crumbly bits, pips, seeds, fibrous parts of fruit, husks or bones 	<p>Pressure from a fork held on its side can be used to 'cut' or break apart or flake this texture into smaller pieces</p>
7 – Regular	<ul style="list-style-type: none"> • Normal everyday foods of various textures that are appropriate for the person • Any method may be used to eat these foods • Food may be hard, crunchy or naturally soft • Includes hard, tough, chewy, fibrous, stringy, dry, crisp, crunchy or crumbly bits • Includes dual texture foods 	Not Applicable

BAPEN MUST Tools

Step 1

BMI score

BMI kg/m ²	Score
>20 (>30 Obese)	= 0
18.5-20	= 1
<18.5	= 2

Step 2

Weight loss score

Unplanned weight loss in past 3-6 months	
%	Score
<5	= 0
5-10	= 1
>10	= 2

Step 3

Acute disease effect score

If patient is acutely ill and there has been or is likely to be no nutritional intake for >5 days
Score 2



If unable to obtain height and weight, see reverse for alternative measurements and use of subjective criteria

Acute disease effect is unlikely to apply outside hospital. See 'MUST' Explanatory Booklet for further information

Step 4

Overall risk of malnutrition

Add Scores together to calculate overall risk of malnutrition
Score 0 Low Risk Score 1 Medium Risk Score 2 or more High Risk

Step 5

Management guidelines

0

Low Risk

Routine clinical care

- Repeat screening
Hospital – weekly
Care Homes – monthly
Community – annually for special groups e.g. those >75 yrs

1

Medium Risk

Observe

- Document dietary intake for 3 days
- If adequate – little concern and repeat screening
 - Hospital – weekly
 - Care Home – at least monthly
 - Community – at least every 2-3 months
- If inadequate – clinical concern – follow local policy, set goals, improve and increase overall nutritional intake, monitor and review care plan regularly

2 or more High Risk

Treat*

- Refer to dietitian, Nutritional Support Team or implement local policy
- Set goals, improve and increase overall nutritional intake
- Monitor and review care plan
Hospital – weekly
Care Home – monthly
Community – monthly
- * Unless detrimental or no benefit is expected from nutritional support e.g. imminent death.

All risk categories:

- Treat underlying condition and provide help and advice on food choices, eating and drinking when necessary.
- Record malnutrition risk category.
- Record need for special diets and follow local policy.

Obesity:

- Record presence of obesity. For those with underlying conditions, these are generally controlled before the treatment of obesity.

Re-assess subjects identified at risk as they move through care settings

See The 'MUST' Explanatory Booklet for further details and The 'MUST' Report for supporting evidence.

© BAPEN

Step 1 – BMI score (& BMI)

Height (feet and inches)

	4'9½	4'10½	4'11	5'0	5'0½	5'1½	5'2	5'3	5'4	5'4½	5'5½	5'6	5'7	5'7½	5'8½	5'9½	5'10	5'11	5'11½	6'0½	6'1	6'2	6'3	6'3½	6'4½	
Weight (kg)	47	48	44	43	42	41	40	39	38	37	38	35	35	34	33	32	32	31	30	30	29	28	28	27	27	15.10
99	48	45	44	43	42	41	40	39	38	37	38	35	34	33	33	32	31	31	30	29	28	28	27	27	28	15.8
98	48	45	44	42	41	40	39	38	37	38	35	34	34	33	33	32	32	31	30	30	29	28	28	27	28	15.6
97	48	44	43	42	41	40	39	38	37	38	35	34	34	33	32	32	31	31	30	30	29	28	28	27	28	15.4
96	45	44	43	42	40	39	38	37	38	35	34	34	33	33	32	32	31	31	30	30	29	28	28	27	28	15.2
95	45	43	42	41	40	39	38	37	38	35	34	34	33	33	32	32	31	31	30	30	29	28	28	27	28	14.13
94	44	43	42	41	40	39	38	37	38	35	34	33	33	32	32	31	31	30	30	29	28	28	27	28	25	14.11
93	44	42	41	40	39	38	37	38	35	34	34	33	33	32	31	31	30	29	28	28	27	28	28	25	25	14.9
92	43	42	41	40	39	38	37	38	35	34	33	33	32	32	31	31	30	30	29	28	28	27	28	26	25	24
91	43	42	40	39	38	37	38	35	34	33	33	32	32	31	31	30	30	29	28	28	27	28	26	25	25	24
90	42	41	40	39	38	37	38	35	34	33	33	32	32	31	30	30	29	28	28	27	27	28	26	25	24	24
89	42	41	40	39	38	37	38	35	34	33	33	32	32	31	30	29	28	27	27	28	26	25	25	24	24	24
88	41	40	39	38	37	36	35	34	34	33	32	31	30	30	29	28	27	27	28	26	25	25	24	24	23	23.12
87	41	40	39	38	37	36	35	34	33	32	32	31	30	30	29	28	27	27	28	26	25	25	24	24	23	23.10
86	40	39	38	37	36	35	34	34	33	32	31	30	30	29	28	27	27	28	26	25	25	24	24	23	23	23.8
85	40	39	38	37	36	35	34	33	32	32	31	30	29	28	28	27	27	28	26	25	25	24	24	23	23	23.5
84	38	37	36	35	35	34	33	32	31	30	30	29	28	28	27	27	28	26	25	25	24	24	23	23	22	13.3
83	38	37	36	35	34	33	32	32	31	30	29	28	27	27	28	26	25	25	24	23	23	22	22	21	21	13.1
82	38	37	36	35	34	33	32	32	31	30	29	28	27	27	28	26	25	25	24	24	23	22	22	21	21	12.13
81	38	37	36	35	34	33	32	32	31	30	29	28	27	27	28	26	25	24	24	23	22	22	21	21	21	12.11
80	38	37	36	35	34	33	32	31	30	29	28	27	27	28	26	25	24	24	23	23	22	22	21	21	21	12.8
79	37	36	35	34	33	32	32	31	30	29	28	28	27	27	28	26	25	24	24	23	23	22	22	21	21	12.6
78	37	36	35	34	33	32	31	30	29	28	28	27	27	28	26	25	25	24	24	23	22	22	21	21	21	12.4
77	36	35	34	33	32	31	30	29	28	28	27	27	28	26	25	24	24	23	22	22	21	21	20	20	20	12.2
76	36	35	34	33	32	31	30	29	28	28	27	26	26	25	25	24	23	22	22	22	21	21	20	20	20	12.0
75	35	34	33	32	32	31	30	29	28	28	27	27	28	26	25	24	24	23	22	22	21	21	20	20	20	11.11
74	35	34	33	32	31	30	29	28	28	27	28	26	25	24	24	23	23	22	22	21	21	20	20	20	20	11.9
73	34	33	32	32	31	30	29	28	28	27	28	26	25	25	24	24	23	23	22	22	21	21	20	20	20	11.7
72	34	33	32	31	30	30	29	28	27	27	28	26	25	24	24	23	23	22	22	21	21	20	20	20	20	11.5
71	33	32	32	31	30	29	28	28	27	28	26	25	25	24	23	23	22	22	21	21	20	20	20	19	19	11.3
70	33	32	31	30	30	29	28	27	27	28	25	25	24	24	23	23	22	22	21	21	20	20	20	20	20	11.0
69	32	32	31	30	29	28	28	27	27	28	25	25	24	24	23	23	22	22	21	21	20	20	20	20	20	10.12
68	32	31	30	29	28	28	27	26	26	25	25	24	24	23	23	22	22	21	21	20	20	19	19	18	18	10.10
67	31	31	30	29	28	28	27	26	26	25	24	24	23	23	22	22	21	21	20	20	19	19	18	18	18	10.8
66	31	30	29	28	28	27	26	25	25	24	23	23	22	22	21	21	20	20	19	19	18	18	18	18	18	10.6
65	30	30	29	28	27	27	26	25	25	24	24	23	22	22	21	21	21	20	20	19	19	18	18	18	17	10.3
64	30	29	28	27	26	25	24	24	23	23	22	22	21	21	20	20	19	19	18	18	17	17	17	17	17	10.1
63	30	29	28	27	26	25	25	24	23	23	22	22	21	21	20	20	19	19	18	18	17	17	17	17	17	9.13
62	29	28	27	26	25	25	24	24	23	22	22	21	21	20	20	20	19	19	18	18	18	17	17	17	17	9.11
61	29	28	27	26	25	24	24	23	23	22	22	21	21	20	20	19	19	18	18	18	17	17	17	17	17	9.8
60	28	27	26	25	25	24	23	23	22	22	21	21	20	20	19	19	18	18	17	17	17	17	17	17	17	9.6
59	28	27	26	25	24	24	23	22	22	21	21	20	20	19	19	18	18	17	17	17	17	17	17	17	17	9.4
58	27	26	25	24	24	23	22	22	21	21	20	20	19	19	18	18	18	18	17	17	17	17	17	17	17	9.2
57	27	26	25	24	23	23	22	22	21	21	20	20	19	19	18	18	18	18	17	17	17	17	17	17	17	9.0
56	26	25	24	24	23	22	22	21	21	20	20	19	19	18	18	17	17	17	17	17	17	17	17	17	17	8.11
55	25	24	23	23	22	22	21	21	20	20	19	19	18	18	17	17	17	17	17	17	17	17	17	17	17	8.9
54	25	24	23	23	22	22	21	21	20	20	19	19	18	18	17	17	17	17	17	17	17	17	17	17	17	8.7
53	25	24	23	22	22	21	21	20	20	19	19	18	18	18	17	17	17	17	17	17	17	17	17	17	17	8.5
52	24	23	23	22	21	21	20	20	19	19	18	18	18	17	17	17	18	18	18	15	15	15	15	14	14	8.3
51	24	23	23	22	22	21	20	20	19	19	18	18	18	17	17	18	18	18	15	15	15	14	14	14	14	8.0
50	23	23	22	21	21	20	20	19	19	18	18	17	17	17	17	18	18	15	15	15	15	14	14	14	13	7.12
49	23	22	22	21																						

Step 2 – Weight loss score

	Score 0 Wt loss < 5%	Score 1 Wt loss 5 - 10%	Score 2 Wt loss > 10%		Score 0 Wt loss < 5%	Score 1 Wt loss 5 - 10%	Score 2 Wt loss > 10%
Weight loss in last 3 to 6 months							
kg	Less than (kg)	Between (kg)	More than (kg)		kg	Less than (kg)	Between (kg)
30	1.6	1.6 - 3.3	3.3		65	3.4	3.4 - 7.2
31	1.6	1.6 - 3.4	3.4		66	3.5	3.5 - 7.3
32	1.7	1.7 - 3.6	3.6		67	3.5	3.5 - 7.4
33	1.7	1.7 - 3.7	3.7		68	3.6	3.6 - 7.6
34	1.8	1.8 - 3.8	3.8		69	3.6	3.6 - 7.7
35	1.8	1.8 - 3.9	3.9		70	3.7	3.7 - 7.8
36	1.9	1.9 - 4.0	4.0		71	3.7	3.7 - 7.9
37	1.9	1.9 - 4.1	4.1		72	3.8	3.8 - 8.0
38	2.0	2.0 - 4.2	4.2		73	3.8	3.8 - 8.1
39	2.1	2.1 - 4.3	4.3		74	3.9	3.9 - 8.2
40	2.1	2.1 - 4.4	4.4		75	3.9	3.9 - 8.3
41	2.2	2.2 - 4.6	4.6		76	4.0	4.0 - 8.4
42	2.2	2.2 - 4.7	4.7		77	4.1	4.1 - 8.6
43	2.3	2.3 - 4.8	4.8		78	4.1	4.1 - 8.6
44	2.3	2.3 - 4.9	4.9		79	4.2	4.2 - 8.7
45	2.4	2.4 - 5.0	5.0		80	4.2	4.2 - 8.9
46	2.4	2.4 - 5.1	5.1		81	4.3	4.3 - 9.0
47	2.5	2.5 - 5.2	5.2		82	4.3	4.3 - 9.1
48	2.5	2.5 - 5.3	5.3		83	4.4	4.4 - 9.2
49	2.6	2.6 - 5.4	5.4		84	4.4	4.4 - 9.3
50	2.6	2.6 - 5.6	5.6		85	4.5	4.5 - 9.4
51	2.7	2.7 - 5.7	5.7		86	4.5	4.5 - 9.6
52	2.7	2.7 - 5.8	5.8		87	4.6	4.6 - 9.7
53	2.8	2.8 - 5.9	5.9		88	4.6	4.6 - 9.8
54	2.8	2.8 - 6.0	6.0		89	4.7	4.7 - 9.9
55	2.9	2.9 - 6.1	6.1		90	4.7	4.7 - 10.0
56	2.9	2.9 - 6.2	6.2		91	4.8	4.8 - 10.1
57	3.0	3.0 - 6.3	6.3		92	4.8	4.8 - 10.2
58	3.1	3.1 - 6.4	6.4		93	4.9	4.9 - 10.3
59	3.1	3.1 - 6.6	6.6		94	4.9	4.9 - 10.4
60	3.2	3.2 - 6.7	6.7		95	5.0	5.0 - 10.6
61	3.2	3.2 - 6.8	6.8		96	5.1	5.1 - 10.7
62	3.3	3.3 - 6.9	6.9		97	5.1	5.1 - 10.8
63	3.3	3.3 - 7.0	7.0		98	5.2	5.2 - 10.9
64	3.4	3.4 - 7.1	7.1		99	5.2	5.2 - 11.0

Alternative measurements: instructions and tables

If height cannot be obtained, use length of forearm (ulna) to calculate height using tables below.

(See *The 'MUST' Explanatory Booklet* for details of other alternative measurements (knee height and demispan) that can also be used to estimate height).


Estimating height from ulna length



Measure between the point of the elbow (olecranon process) and the midpoint of the prominent bone of the wrist (styloid process) (left side if possible).


Height (m)	men (<65 years)	1.94	1.93	1.91	1.89	1.87	1.85	1.84	1.82	1.80	1.78	1.76	1.75	1.73	1.71
	men (≥65 years)	1.87	1.86	1.84	1.82	1.81	1.79	1.78	1.76	1.75	1.73	1.71	1.70	1.68	1.67
	Ulna length (cm)	32.0	31.5	31.0	30.5	30.0	29.5	29.0	28.5	28.0	27.5	27.0	26.5	26.0	25.5
Height (m)	Women (<65 years)	1.84	1.83	1.81	1.80	1.79	1.77	1.76	1.75	1.73	1.72	1.70	1.69	1.68	1.66
	Women (≥65 years)	1.84	1.83	1.81	1.79	1.78	1.76	1.75	1.73	1.71	1.70	1.68	1.66	1.65	1.63
Height (m)	men (<65 years)	1.69	1.67	1.66	1.64	1.62	1.60	1.58	1.57	1.55	1.53	1.51	1.49	1.48	1.46
	men (≥65 years)	1.65	1.63	1.62	1.60	1.59	1.57	1.56	1.54	1.52	1.51	1.49	1.48	1.46	1.45
	Ulna length (cm)	25.0	24.5	24.0	23.5	23.0	22.5	22.0	21.5	21.0	20.5	20.0	19.5	19.0	18.5
Height (m)	Women (<65 years)	1.65	1.63	1.62	1.61	1.59	1.58	1.56	1.55	1.54	1.52	1.51	1.50	1.48	1.47
	Women (≥65 years)	1.61	1.60	1.58	1.56	1.55	1.53	1.52	1.50	1.48	1.47	1.45	1.44	1.42	1.40

Estimating BMI category from mid upper arm circumference (MUAC)



The subject's left arm should be bent at the elbow at a 90 degree angle, with the upper arm held parallel to the side of the body. Measure the distance between the bony protrusion on the shoulder (acromion) and the point of the elbow (olecranon process). Mark the mid-point.

Ask the subject to let arm hang loose and measure around the upper arm at the mid-point, making sure that the tape measure is snug but not tight.



If MUAC is <23.5 cm, BMI is likely to be <20 kg/m².

If MUAC is >32.0 cm, BMI is likely to be >30 kg/m².

The use of MUAC provides a general indication of BMI and is not designed to generate an actual score for use with 'MUST'. For further information on use of MUAC please refer to *The 'MUST' Explanatory Booklet*.

Useful Web Links

- MUST Calculator: ['MUST' Calculator | BAPEN](#)
- MUST Tool Kit for downloadable charts: [The 'MUST' Toolkit | BAPEN](#)
- Texture modified diet criteria and further information:
[V2DetailedDefnEnglish31july2019.pdf](#) [The IDDSI Standard - Standards - IDDSI](#)
- Food Fortification Video: [\(9396\) Food Fortification - Devised by the North Derbyshire Dietitians - YouTube](#)
- Thickening Fluids Video: [How to thicken fluids: levels 1 to 4 - Chesterfield Royal Hospital NHS Foundation Trust](#)
- BDA Nutrition and Hydration Digest [Care Home Digest - BDA](#)
- Derbyshire Medicines Management Vitamin D information:
[Position Statement for Vit D.pdf](#)
- Derbyshire Medicines Management of Undernutrition in Adults: [Management of undernutrition in adults](#)
- Big Nutrition for Small Appetite Leaflets: [Big Nutrition for Small Appetites](#)
- Further information on CQC Guideline 14 – Meeting Nutritional and Hydration Needs: [Regulation 14: Meeting nutritional and hydration needs - Care Quality Commission](#)
- Malnutrition Pathway Protein resource: [Why it is important and where to find it](#)