


Derbyshire commissioning guidance for the treatment of Psoriatic Arthritis
October 2018



Dosing schedule

Biologic	NICE TA	Loading dose	Maintenance dose	Response measured	
Adalimumab (SC)	TA199	40mg every 2 weeks	NA	12 weeks	Monoclonal antibody that specifically binds to TNF
Etanercept (SC)	TA199	50mg once weekly	NA	12 weeks	Recombinant human TNF receptor fusion protein.
Golimumab (SC)	TA220	50mg every month >100kg in body weight, 100mg every month after 3-4 initial doses.	NA	12 weeks	Monoclonal antibody that prevents the binding of TNF to its receptors.
Certolizumab (SC)	TA445	Week 0,2 & 4 - 400mg	200mg every 2 weeks or 400mg every 4 weeks	12 weeks	Recombinant humanised antibody Fab' fragment against TNF alpha
Secukinumab (SC)	TA445	For patients with concomitant moderate to severe plaque psoriasis: Week 0,1,2 & 3 – 300mg For other patients: Week 0,1,2 & 3 – 150mg	Week 4 – 300mg & then continue every month. Followed by monthly maintenance dosing 150mg starting at week 4.	16 weeks	Secukinumab is a high-affinity, fully human monoclonal antibody that binds to and neutralises interleukin-17A,
Ixekizumab (SC)	TA537	Week 0 – 160mg For patients with moderate to severe plaque psoriasis Week 0 – 160mg Week 2 - 80mg Week 4 – 80mg Week 6 – 80mg Week 8 – 80mg Week 10 – 80mg Week 12 – 80mg	Every 4 weeks Every 4 weeks thereafter.	16 weeks	Ixekizumab) is an antibody that inhibits IL-17A (interleukin-17A, a pro-inflammatory cytokine).
Ustekinumab (SC)	TA340	Week 0 & 4 - 45mg or >100kg in body weight – 90mg	Every 12 weeks thereafter.	24 weeks	Ustekinumab is a fully human monoclonal antibody that targets interleukin-12 (IL-12) and IL-23
Intravenous infusion					
Infliximab (IV)	TA199	Week 0, 2 & 6 - 5mg/kg IV	5mg/kg IV every 8 weeks thereafter	12 weeks	Chimeric monoclonal antibody, with high affinity to TNF.
Oral preparations					
Apremilast (PO)	TA433	<ul style="list-style-type: none"> Day 1 - 10mg am Day 2 - 10mg am & pm Day 3 - 10mg am, 20mg pm Day 4 - 20mg am & pm Day 5 - 20mg am & 30mg pm 	Day 6 and thereafter - 30mg am & pm	16 weeks	Apremilast is an oral small-molecule inhibitor of phosphodiesterase 4 (PDE4), works intracellularly to modulate a network of pro-inflammatory and anti-inflammatory mediators

Date Reviewed: October 2018

Review Date: September 2020