

CLINICAL POLICY ADVISORY GROUP (CPAG)

Spinal Decompression for Sciatica Policy

Statement

Derby and Derbyshire CCG (DDCCG) has deemed that spinal decompression is not routinely commissioned for patients with sciatica, unless criteria 1 **and** 2 are met:

1. Non-surgical treatment has not improved pain or function,
AND
2. Radiological findings are consistent with sciatic symptoms.

Exception

Lumbar discectomy is a type of spinal decompression intervention that is an exception to the DDCCG Spinal Decompression for Sciatica Policy as it requires the additional time frame criteria of symptoms lasting three months (except in severe cases). Therefore, lumbar discectomy is not routinely commissioned for patients with sciatica, unless criteria 1, 2 and 3 are **ALL** met:

1. Non-surgical treatment has not improved pain or function,
AND
2. Radiological findings are consistent with sciatic symptoms
AND
3. Symptoms have lasted for 3 months or longer (except in severe cases)

This policy applies to patients aged 16 years and over.

This commissioning statement is aligned with the [Evidence-Based Interventions List 2 Guidance](#).

The CCG's commissioning intentions will be reviewed periodically. This is to ensure affordability against other services commissioned by the CCG.

1. Background Information

Spinal decompression surgery is a type of surgery used to treat compressed nerves in the lower (lumbar) spine while maintaining as much of the strength and flexibility of the spine as possible. Spinal decompression surgery is only recommended when non-surgical treatments haven't helped.

Depending on the specific reason for having surgery, a number of different types of spinal decompression procedures may need to be carried out during the operation to treat the compressed nerves.

Examples of the main procedures used include:

- laminectomy – where a section of bone is removed from one of your vertebrae (spinal bones) to relieve pressure on the affected nerve
- discectomy – where a section of a damaged disc is removed

2. Recommendations

Spinal decompression is not routinely commissioned for patients with sciatica, unless criteria 1 **AND** 2 are met:

1. Non-surgical treatment has not improved pain or function,
AND
2. Radiological findings are consistent with sciatic symptoms

Exception

Lumbar discectomy is a type of spinal decompression intervention that is an exception to the DDCCG Spinal Decompression for Sciatica Policy as it requires the additional time frame criteria of symptoms lasting three months (except in severe cases). Therefore, lumbar discectomy is not routinely commissioned for patients with sciatica, unless criteria 1, 2 and 3 are **ALL** met:

1. Non-surgical treatment has not improved pain or function,
AND
2. Radiological findings are consistent with sciatic symptoms
AND
3. Symptoms have lasted for 3 months or longer (except in severe cases)

This policy applies to patients aged 16 years and over.

This commissioning statement is aligned with the [Evidence-Based Interventions List 2 Guidance](#)

Exceptional Circumstances

DDCCG will consider individual cases for funding outside this commissioning statement in accordance with their [Individual Funding Request \(IFR\) Policy](#) which sets out a decision-making framework for determining these cases.

For an IFR request to be considered, it must be demonstrated that the patient fulfils the strict criteria for exceptionality. It should be noted that the criteria for exceptionality is very unlikely

to be satisfied if an individual is part of an identifiable cohort of patients, who at the same disease stage would derive similar benefit from the intervention.

3. Rationale for Recommendations

There remains a reasonable body of evidence to show that in carefully selected patients, lumbar discectomy may lead to a greater and quicker improvement in pain scores than in non-operatively treated patients.

In other studies however, because of the irreversible degenerative changes, surgery has not shown a benefit over non-operative treatment in mid and long-term follow-up.

Lengthy periods of ineffective non-operative care may prompt repeated emergency department attendances, issues with chronic pain, significant neurological dysfunction and time off work.

4. Useful Resources

- Lumbar decompression surgery, NHS, last reviewed 21/07/18, accessed 09/06/21, <https://www.nhs.uk/conditions/lumbar-decompression-surgery/>

5. Reference

- NICE Guidance [NG59], Low back pain and sciatica in over 16s: assessment and management, November 2016, accessed 31/12/19, <https://www.nice.org.uk/guidance/ng59>
- Evidence-Based Interventions - List 2 Guidance, Academy of Medical Colleges, November 2020, accessed 09/06/21, https://www.aomrc.org.uk/wp-content/uploads/2020/12/EBI_list2_guidance_150321.pdf
- Lumbar decompression surgery, NHS, last reviewed 21/07/18, accessed 09/06/21, <https://www.nhs.uk/conditions/lumbar-decompression-surgery/>

6. Appendices

Appendix 1- Consultation

All relevant providers/stakeholders will be consulted via a named link consultant/specialist. Views expressed should be representative of the provider/stakeholder organisation. CPAG will consider all views to inform a consensus decision, noting that sometimes individual views and opinions will differ.

Consultee	Date
Consultant Trauma, Orthopaedic & Spinal Surgeon, UHDB	May 2021

Clinical Policy Advisory Group	June 2021
Clinical Lay Commissioning Committee	July 2021

Appendix 2- Document Update

Document Update	Date Updated
Version 2 - Derby and Derbyshire CCG Position Statement for Spinal Decompression reformatted into a clinical policy format. Addition of the intervention lumbar discectomy to the policy under the new 'Exception' section of the policy.	June 2021