



Publications gateway number: GOV-14998

Live attenuated influenza vaccine nasal spray suspension (LAIV) Patient Group Direction (PGD)

This PGD is for the supply and administration, or supply only, of live attenuated influenza vaccine (LAIV) nasal spray suspension (Fluenz[®] Tetra) to children and adolescents from 2 years to under 18 years of age in accordance with the national flu immunisation programme.

This PGD is for the supply and administration, or supply only, of LAIV by registered healthcare practitioners identified in [Section 3](#), subject to any limitations to authorisation detailed in [Section 2](#).

Reference no: LAIV PGD
Version no: v13.00
Valid from: 1 September 2023
Review date: 1 April 2024
Expiry date: 1 April 2024

The UK Health Security Agency (UKHSA) has developed this PGD to facilitate the delivery of publicly funded immunisations in England in line with national recommendations.

Those using this PGD must ensure that it is organisationally authorised and signed in Section 2 by an appropriate authorising person, relating to the class of person by whom the product is to be supplied, in accordance with Human Medicines Regulations 2012 (HMR2012)¹. **The PGD is not legal or valid without signed authorisation in accordance with [HMR2012 Schedule 16 Part 2](#).**

Authorising organisations must not alter, amend or add to the clinical content of this document (sections 4, 5 and 6); such action will invalidate the clinical sign-off with which it is provided. In addition authorising organisations must not alter section 3 'Characteristics of staff'. Only sections 2 and 7 can be amended within the designated editable fields provided.

Operation of this PGD is the responsibility of commissioners and service providers. The final authorised copy of this PGD should be kept by the authorising organisation completing Section 2 for 8 years after the PGD expires if the PGD relates to adults only and for 25 years after the PGD expires if the PGD relates to children only, or adults and children. Provider organisations adopting authorised versions of this PGD should also retain copies for the periods specified above.

Individual practitioners must be authorised by name, under the current version of this PGD before working according to it.

Practitioners and organisations must check that they are using the current version of the PGD. Amendments may become necessary prior to the published expiry date. Current versions of UKHSA PGD templates for authorisation can be found from:

[Immunisation patient group direction \(PGD\) templates](#)

Any concerns regarding the content of this PGD should be addressed to:
immunisation@ukhsa.gov.uk

Enquiries relating to the availability of organisationally authorised PGDs and subsequent versions of this PGD should be directed to: [Vaccination and Screening Programmes, NHS England – Midlands, responsible for your area:

¹ This includes any relevant amendments to legislation.

East: england.emids-imms@nhs.net

- Derby and Derbyshire
- Lincolnshire
- Leicester, Leicestershire and Rutland
- Northamptonshire
- Nottingham and Nottinghamshire

West: england.wmid-imms@nhs.net

- Herefordshire and Worcestershire
- Birmingham and Solihull
- Staffordshire and Stoke-on-Trent
- Shropshire, Telford and Wrekin
- Black Country
- Coventry and Warwickshire

Change history




Version number	Change details ²	Date
Final version	New PHE Fluenz PGD	1 September 2013
Final version – revised	See earlier version of this PGD for change details.	9 September 2013
V2.00	See earlier version of this PGD for change details.	11 August 2015
V3.00	See earlier version of this PGD for change details.	20 October 2015
V4.00	See earlier version of this PGD for change details.	22 June 2016
V5.00	See earlier version of this PGD for change details.	4 July 2017
V6.00	See earlier version of this PGD for change details.	17 August 2017
V7.00	See earlier version of this PGD for change details.	8 June 2018
V8.00	<p>PHE LAIV PGD amended to:</p> <ul style="list-style-type: none"> include the 2019 to 2020 influenza programme eligible cohorts, with the addition of children of appropriate age for school year 6 remove the exclusion of individuals on high dose inhaled corticosteroids and replace with the exclusion of individuals who require oral steroid for the maintenance of asthma control or have previously required intensive care for an asthma exacerbation, in accordance with updated recommendations from JCVI and in Chapter 19 of the Green Book include reference to the Directed Enhanced Service and offer to morbidly obese adults from 16 years of age include minor rewording, layout and formatting changes to remove duplication and for clarity and consistency with other PHE PGD templates 	8 May 2019
V9.00	<p>PHE LAIV PGD amended to:</p> <ul style="list-style-type: none"> extend the characteristics of staff to include all registered practitioners legally able to work under PGD include the 2020 to 2021 influenza programme eligible DOB cohorts and household contacts of those on the COVID-19 Shielded Patient List include minor rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	16 July 2020
V10.00	<p>PHE LAIV PGD amended to:</p> <ul style="list-style-type: none"> include the 2021 to 2022 influenza programme eligible cohorts 	28 July 2021
V11.00	<p>LAIV PGD amended to:</p> <ul style="list-style-type: none"> include the 2022 to 2023 influenza vaccination programme eligible cohorts update organisation from PHE to the UKHSA 	9 August 2022

² A summary of the changes between superseded versions may be found in more detail by referring to the Change History in the relevant earlier versions of this PGD.

V12.00	<p>LAIV PGD amended to:</p> <ul style="list-style-type: none"> • update to include the 2023 to 2024 influenza vaccination programme eligible cohorts • align the very severely immunocompromised statement to Green Book in the inclusion criteria • add children aged 2 years to less than 9 years who are household contacts of immunocompromised individuals in dose section • add additional information about use of the nasal spray in the route and administration section • update the owner of the technical memorandum from Department of Health to NHSE in the disposal section • move headache to less common adverse reaction in line with the SPC in identification of adverse reactions • add use of salicylate caution in patient advice section • add how to obtain accessible information in the written information given to patient section • updated the references 	29 June 2023
V13.00	<p>LAIV PGD amended to:</p> <ul style="list-style-type: none"> • add secondary school Years 7-11 to criteria for inclusion section 	4 July 2023

1. PGD development

This PGD has been developed by the following health professionals on behalf of the UKHSA:

Developed by:	Name	Signature	Date
Pharmacist (Lead Author)	Suki Hunjunt Lead Pharmacist Immunisation Services, Immunisation and Vaccine Preventable Diseases Division, UKHSA		10 July 2023
Doctor	Jamie Lopez-Bernal Consultant Epidemiologist, Immunisation and Vaccine Preventable Diseases Division, UKHSA		10 July 2023
Registered Nurse (Chair of Expert Panel)	David Green Nurse Consultant for Immunisation, Immunisation and Vaccine Preventable Diseases Division, UKHSA		10 July 2023

This PGD has been peer reviewed by the UKHSA Immunisations PGD Expert Panel in accordance with the UKHSA PGD Policy. It has been ratified by the UKHSA Medicines Governance Group.

Expert Panel

Nicholas Aigbogun	Consultant in Communicable Disease Control, Yorkshire and Humber Health Protection Team, UKHSA
Gayatri Amrithalingham	Consultant Epidemiologist, Immunisation and Vaccine Preventable Diseases Division, UKHSA
Alison Campbell	Screening and Immunisation Coordinator, Public Health Commissioning NHS England (NHSE) Midlands
Sarah Dermont	Clinical Project Coordinator and Registered Midwife, NHS Infectious Diseases in Pregnancy Screening Programme, NHSE
Rosie Furner	Pharmacist - Medicines Governance, Specialist Pharmacist Services (SPS)
Ed Gardner	Advanced Paramedic Practitioner/Emergency Care Practitioner, Medicines Manager, Proactive Care Lead
Jacqueline Lamberty	Lead Pharmacist, Medicines Governance, UKHSA
Michelle Jones	Principal Medicines Optimisation Pharmacist, Bristol North Somerset and South Gloucestershire Integrated Care Board
Shamez Ladhani	Paediatric Infectious Disease Consultant, UKHSA
Elizabeth Lockett	Senior Screening & Immunisation Manager NHSE South West
Vanessa MacGregor	Consultant in Communicable Disease Control, East Midlands Health Protection Team, UKHSA
Suzanna McDonald	National Programme Lead for Influenza, UKHSA
Lesley McFarlane	Lead Immunisation Nurse Specialist Immunisation and Vaccine Preventable Diseases Division, UKHSA
Nicola Philbin	Screening and Immunisation Manager, Vaccination and screening programmes – Public Health Commissioning NHSE Midlands

Conall Watson	Consultant Epidemiologist – influenza and seasonal respiratory viruses, UKHSA
Tushar Shah	Lead Pharmacy Adviser, NHSE London


2. Organisational authorisations

The PGD is not legally valid until it has had the relevant organisational authorisation.

It is the responsibility of the organisation that has legal authority to authorise the PGD, to ensure that all legal and governance requirements are met. The authorising body accepts governance responsibility for the appropriate use of the PGD.

NHSE Midlands authorises this PGD for use by the services or providers listed below:

Authorised for use by the following organisations and/or services
<p>Primary care services and/or all organisations commissioned or contracted by NHS England – Midlands to provide immunisation services in:</p> <ul style="list-style-type: none"> • Derby and Derbyshire • Lincolnshire • Leicester, Leicestershire, and Rutland • Northamptonshire • Nottingham and Nottinghamshire • Herefordshire and Worcestershire • Birmingham and Solihull • Staffordshire and Stoke-on-Trent • Shropshire, Telford, and Wrekin • Black Country • Coventry and Warwickshire
Limitations to authorisation
None

Organisational approval (legal requirement)			
Role	Name	Sign	Date
Director Primary Care and Public Health Commissioning – Midlands NHS England.	Trish Thompson		01.08.23

Additional signatories according to locally agreed policy			
Role	Name	Sign	Date

Local enquiries regarding the use of this PGD may be directed to: Vaccination and Screening Programmes, NHS England – Midlands, responsible for your area:

East: england.emids-imms@nhs.net

- Derby and Derbyshire
- Lincolnshire
- Leicester, Leicestershire and Rutland
- Northamptonshire
- Nottingham and Nottinghamshire

West: england.wmid-imms@nhs.net

- Herefordshire and Worcestershire
- Birmingham and Solihull
- Staffordshire and Stoke-on-Trent
- Shropshire, Telford and Wrekin
- Black Country
- Coventry and Warwickshire

[Section 7](#) provides a practitioner authorisation sheet. Individual practitioners must be authorised by name to work to this PGD. Alternative practitioner authorisation sheets may be used where appropriate in accordance with local policy but this should be an individual agreement or a multiple practitioner authorisation sheet as included at the end of this PGD.

3. Characteristics of Staff


<p>Qualifications and professional registration required</p>	<p>Practitioners must only work under this PGD where they are competent to do so. Practitioners working to this PGD must also be one of the following registered professionals who can legally supply and administer under a PGD (see Patient Group Directions: who can administer them):</p> <ul style="list-style-type: none"> • nurses and midwives currently registered with the Nursing and Midwifery Council (NMC) • pharmacists currently registered with the General Pharmaceutical Council (GPhC) (Note: This PGD is not relevant to the national community pharmacy seasonal influenza vaccination advanced service nor privately provided community pharmacy services) • chiropodists/podiatrists, dieticians, occupational therapists, orthoptists, orthotists/prosthetists, paramedics, physiotherapists, radiographers and speech and language therapists currently registered with the Health and Care Professions Council (HCPC) • dental hygienists and dental therapists registered with the General Dental Council • optometrists registered with the General Optical Council <p>Practitioners must also fulfil all the Additional requirements.</p> <p>Check Section 2 Limitations to authorisation to confirm whether all the registered practitioners listed above have organisational authorisation to work under this PGD.</p>
<p>Additional requirements</p>	<p>Additionally, practitioners:</p> <ul style="list-style-type: none"> • must be authorised by name as an approved practitioner under the current terms of this PGD before working to it • must have undertaken appropriate training for working under PGDs for supply/administration of medicines • must be competent in the use of PGDs (see NICE Competency framework for health professionals using PGDs) • must be familiar with the vaccine product and alert to changes in the Summary of Product Characteristics (SPC), Immunisation Against Infectious Disease ('The Green Book'), and national and local immunisation programmes • must have undertaken training appropriate to this PGD as required by local policy and in line with the National Minimum Standards and Core Curriculum for Immunisation Training. For further information see Flu immunisation training recommendations • must be competent to undertake immunisation and to discuss issues related to immunisation • must be competent in the handling and storage of vaccines, and management of the cold chain • must be competent in the recognition and management of anaphylaxis • must have access to the PGD and associated online resources • should fulfil any additional requirements defined by local policy <p>The individual practitioner must be authorised by name, under the current version of this PGD before working according to it.</p>
<p>Continued training requirements</p> <p>Continued over page</p>	<p>Practitioners must ensure they are up to date with relevant issues and clinical skills relating to immunisation and management of anaphylaxis, with evidence of appropriate Continued Professional Development (CPD).</p> <p>Practitioners should be constantly alert to any subsequent recommendations from the UKHSA and/or NHSE and other sources of medicines information. Note: The most current national recommendations should be followed but a Patient Specific Direction (PSD) may be required to administer the vaccine</p>

Continued training requirements (continued)	in line with updated recommendations that are outside the criteria specified in this PGD.
---	---

4. Clinical condition or situation to which this PGD applies.

<p>Clinical condition or situation to which this PGD applies</p>	<p>LAIV is indicated for the active immunisation of children and adolescents from 2 years to under 18 years of age for the prevention of influenza infection, in line with the recommendations given in Chapter 19 of Immunisation Against Infectious Disease: 'The Green Book' and annual flu letter(s).</p>
<p>Criteria for inclusion</p> <p>Continued over page Criteria for inclusion (continued)</p>	<p>Individuals eligible for vaccination with LAIV in accordance with national recommendations for the 2023 to 2024 influenza season including:</p> <ul style="list-style-type: none"> • all those aged 2 or 3 years on 31 August 2023 (with a date of birth on or after 1 September 2019 and on or before 31 August 2021) • all primary school-aged children in reception to Year 6 (aged 4 to 10 years old on 31 August 2023) including home-schooled and other children not in mainstream education <ul style="list-style-type: none"> ○ some school aged children might be outside of the age ranges outlined in the above paragraphs (for example, if a child has been accelerated or held back a year). It is acceptable to offer and deliver influenza immunisation to these children with their class peers under this PGD • secondary school-aged children in Years 7-11 including home-schooled and other children not in mainstream education • children and adolescents from 2 years to under 18 years of age who are in a clinical risk group category listed in Chapter 19 of the Green Book such as: <ul style="list-style-type: none"> ○ chronic (long-term) respiratory disease, such as asthma (that requires continuous or repeated use of inhaled or systemic steroids or with previous exacerbations requiring hospital admission, but see also criteria for exclusion below), bronchitis or cystic fibrosis ○ chronic heart disease and vascular disease, such as heart failure ○ chronic kidney disease at stage 3, 4 or 5 ○ chronic liver disease ○ chronic neurological disease, such as cerebral palsy or motor neurone disease ○ learning disability ○ diabetes and adrenal insufficiency ○ asplenia or dysfunction of the spleen ○ a weakened immune system due to disease (such as HIV/AIDS) or treatment (such as cancer treatment) ○ morbidly obese adults (aged from 16 years) with a BMI $\geq 40\text{kg/m}^2$ • children and adolescents from 2 years to under 18 years of age who are close household contacts of immunocompromised individuals. This would also include individuals who expect to share living accommodation on most days over the winter with immunocompromised individuals and therefore for whom continuing close contact is unavoidable. This may include carers. Note: close contacts (example household members or carers) of very severely immunocompromised individuals, for example bone marrow transplant patients requiring isolation, should receive inactivated influenza vaccine and not LAIV, see Inactivated Influenza PGD • individuals, from 16 years to under 18 years of age, who are in receipt of a carer's allowance, or those who are the main carer of an older or disabled person whose welfare may be at risk if the carer falls ill • frontline staff from 16 years to under 18 years of age without employer led occupational health schemes employed: <ul style="list-style-type: none"> ○ by a registered residential care or nursing home or registered domiciliary care provider, who are directly involved in the care of vulnerable individuals who are at increased risk from exposure to influenza

	<p>protection in this high-risk group (see the UKHSA Inactivated Influenza PGD).</p> <p>Seek appropriate advice from the local Screening and Immunisation Team, local Health Protection Team or individual's clinician as required.</p> <p>The risk to the individual of not being immunised must be taken into account.</p> <p>Document the reason for exclusion and any action taken in the individual's clinical records.</p> <p>Inform or refer to the GP or a prescriber as appropriate.</p>
Action to be taken if the patient or carer declines treatment	<p>Informed consent, from the individual or a person legally able to act on the individual's behalf, must be obtained for each administration. For further information on consent see Chapter 2 of the Green Book.</p> <p>Advise the individual/parent/carer about the protective effects of the vaccine, the risks of infection and potential complications.</p> <p>If the parent or carer of an eligible child refuses LAIV because of its porcine gelatin content (and they understand that it is the most effective product in the programme), advise the individual/parent/carer they can request an alternative injectable vaccine. UKHSA has procured QIVc for these children. Refer to the UKHSA Inactivated Influenza Vaccine PGD.</p> <p>Document the advice given and decision reached.</p> <p>Inform or refer to the GP or prescriber as appropriate.</p>
Arrangements for referral for medical advice	<p>As per local policy</p>

	<p>The individual can breathe normally during vaccine administration and there is no need to actively inhale or sniff.</p> <p>Administration does not need to be repeated if the individual sneezes or blows their nose immediately following administration.</p> <p>The SPC provides further guidance on administration.</p> <p>Instructions for administration</p>  <table border="1" data-bbox="491 573 1414 779"> <tr> <td data-bbox="491 573 730 779">Remove protective tip cap. Do not remove the dose-divider</td> <td data-bbox="730 573 967 779">With the patient upright, position the applicator and depress as rapidly as possible</td> <td data-bbox="967 573 1193 779">Pinch and remove the dose-divider clip from the plunger</td> <td data-bbox="1193 573 1414 779">Administer the remaining vaccine into the other nostril</td> </tr> </table>	Remove protective tip cap. Do not remove the dose-divider	With the patient upright, position the applicator and depress as rapidly as possible	Pinch and remove the dose-divider clip from the plunger	Administer the remaining vaccine into the other nostril
Remove protective tip cap. Do not remove the dose-divider	With the patient upright, position the applicator and depress as rapidly as possible	Pinch and remove the dose-divider clip from the plunger	Administer the remaining vaccine into the other nostril		
<p>Dose and frequency of administration</p>	<p>Single dose of 0.2ml of LAIV administered as 0.1ml in each nostril.</p> <p>Children in clinical risk groups</p> <p>Children aged 2 years to less than 9 years who are in a clinical risk group category listed in Chapter 19 of the Green Book and who have not received influenza vaccine before, should receive a second dose of LAIV at least 4 weeks after the first dose.</p> <p>Children aged 2 years to less than 9 years who are household contacts of immunocompromised individuals should be vaccinated in accordance with the advice on children in clinical risk groups (see Chapter 19).</p> <p>Second dose of 0.2ml of LAIV administered as 0.1ml in each nostril.</p>				
<p>Duration of treatment</p>	<p>See section on Dose.</p>				
<p>Quantity to be supplied or administered</p>	<p>0.2ml dose to be administered as 0.1ml in each nostril, or 0.2ml of LAIV to be supplied to the individual for immediate self-administration or administration by another person within the clinic setting. Vaccine supplies which are not legally over-labelled for individual use must be administered prior to the individual leaving the immunisation session. Note: The act of administration by anyone other than the registered professional named in Section 7 is outside the remit of this PGD and should only take place in well-defined local circumstances covered by training and local operating protocols.</p> <p>Children aged 2 years to less than 9 years old in a clinical risk category and receiving influenza immunisation for the first time</p> <p>This dose (0.2ml) should be repeated after a 4-week interval.</p>				
<p>Supplies</p> <p>Continued over page</p> <p>Supplies (continued)</p>	<p>LAIV has been purchased centrally for children. These vaccines should be ordered as per the usual mechanisms for the routine childhood immunisation programme.</p> <p>Protocols for the ordering, storage and handling of vaccines should be followed to prevent vaccine wastage (see Green Book Chapter 3).</p>				
<p>Storage</p>	<p>Store at +2°C to +8°C. Store in original packaging in order to protect from light. Do not freeze.</p>				

	<p>Before use, the vaccine may be removed from the cold-chain, without being replaced, for a maximum period of 12 hours at a temperature not above 25°C. If the vaccine has not been used after this 12-hour period, it should be disposed of.</p> <p>In the event of an inadvertent or unavoidable deviation of these conditions vaccine that has been stored outside the conditions stated above should be quarantined and risk assessed for suitability of continued off-label use or appropriate disposal. Refer to Vaccine Incident Guidance.</p>
Disposal	<p>Equipment used for immunisation, including discharged or partially discharged vaccines in an applicator, should be disposed of safely, as medicinally-contaminated clinical waste for incineration, in a yellow UN-approved waste receptacle (this is usually a sharps box), according to local authority arrangements and guidance in the technical memorandum 07-01 (NHSE, 2022).</p>
Drug interactions	<p>There is a potential for influenza antiviral agents to lower the effectiveness of the LAIV. Therefore, influenza antiviral agents and LAIV should not be administered concomitantly.</p> <p>There is a potential for influenza antiviral agents to lower the effectiveness of LAIV. LAIV should be delayed until 48 hours following the cessation of treatment with influenza antiviral agents.</p> <p>Administration of influenza antiviral agents within the 2 weeks following administration of LAIV may adversely affect the effectiveness of the vaccine.</p> <p>Do not administer LAIV to those receiving salicylate therapy (other than topical treatment for localised conditions) and do not use salicylates for 4 weeks after vaccination.</p> <p>LAIV can be given at the same time as other vaccines.</p> <p>Live vaccines which replicate in the mucosa, such as live attenuated influenza vaccine (LAIV) are unlikely to be seriously affected by concomitant COVID-19 vaccination. It is generally better for vaccination to proceed to avoid any further delay in protection and to avoid the risk of the individual not returning for a later appointment (see Chapter19).</p> <p>A detailed list of drug interactions is available in the SPC, which is available from the electronic medicines compendium website.</p>
Identification and management of adverse reactions	<p>The most common adverse reactions observed after administration of LAIV are decreased appetite, nasal congestion, rhinorrhoea, malaise. Less common reactions include myalgia, headache and pyrexia and uncommon reactions include hypersensitivity reactions, epistaxis and rash.</p> <p>A detailed list of adverse reactions is available in the SPC, which is available from the electronic medicines compendium website.</p>
<p>Reporting procedure of adverse reactions</p> <p>Continued over page</p> <p>Reporting procedure of adverse reactions (continued)</p>	<p>Healthcare professionals and individuals/parents/carers are encouraged to report suspected adverse reactions to the Medicines and Healthcare products Regulatory Agency (MHRA) using the Yellow Card reporting scheme or search for MHRA Yellow Card in the Google Play or Apple App Store.</p> <p>Any adverse reaction to the vaccine should be documented in the individual's record and the individual's GP should be informed.</p>

<p>Written information to be given to patient or carer</p>	<p>Manufacturer's packaging is required to include a patient information leaflet (PIL) which should accompany the supply of vaccine under this PGD.</p> <p>If applicable, inform the individual/parent/carer that PIL with large print, Braille or audio CD can be ordered from the manufacturer (see electronic medicines compendium).</p> <p>When LAIV is administered there is no legal requirement to provide the manufacturer's PIL to the individual at the time of administration, although this may be considered good practice.</p> <p>If applicable, inform the individual/parent/carer that PIL with large print, Braille or audio CD can be ordered from the manufacturer (see electronic medicines compendium).</p>
<p>Patient advice and follow up treatment</p>	<p>Inform the individual/parent/carer of possible side effects and their management.</p> <p>The individual/parent/carer should be advised when to seek medical advice in the event of a severe adverse reaction.</p> <p>When applicable, advise the individual/parent/carer when the subsequent dose is due.</p> <p>The individual/parent/carer should be advised not to give acetylsalicylic acid or salicylates (a substance present in many medicines used to relieve pain and lower fever) to the child or adolescent for 4 weeks after vaccination with Fluenz Tetra as there is a risk of Reye's syndrome. However, topical treatment containing acetylsalicylic acid or salicylates for localised conditions can be used.</p> <p>The individual/parent/carer should be informed that LAIV has the theoretical potential for transmission to immunocompromised contacts. Vaccine recipients should attempt to avoid, whenever possible, close association with very severely immunocompromised individuals (such as bone marrow transplant recipients requiring isolation) for 1-2 weeks following vaccination. If the PGD is used for supply only, advise the individual/parent/carer of the process they need to follow for subsequent administration, for instance refer them immediately to an appropriately trained HCSW within the clinic setting.</p> <p>When administration is postponed advise the individual/parent/carer when to return for vaccination.</p>
<p>Special considerations and additional information</p> <p>Continued over page Special considerations and additional information (continued)</p>	<p>As with most vaccines, appropriate medical treatment and supervision should always be readily available in case of an anaphylactic event following the administration of LAIV. Ensure there is immediate access to adrenaline (epinephrine) 1 in 1000 injection and easy access to a telephone.</p> <p>For children under the age of 16 years, those assessed as Gillick competent can self-consent. For further information on consent see Chapter 2 of the Green Book.</p> <p>Minor illnesses without fever or systemic upset are NOT valid reasons to postpone immunisation. If an individual is acutely unwell, immunisation may be postponed until they have fully recovered. This is to avoid confusing the differential diagnosis of any acute illness by wrongly attributing signs or symptoms to adverse effects of the vaccine.</p> <p>LAIV is not contraindicated for use in children or adolescents with stable HIV infection receiving antiretroviral therapy; or who are receiving topical corticosteroids, inhaled corticosteroids, low-dose systemic corticosteroids or those receiving corticosteroids as replacement therapy (such as for</p>

	<p>adrenal insufficiency) or low-dose immunosuppressive therapy. This PGD may be used for these individuals.</p> <p>Individuals with learning disabilities may require reasonable adjustments to support vaccination (see Flu vaccinations: supporting people with learning disabilities). A PSD may be required.</p> <p>LAIV should be offered to eligible children aged from 2 years to less than 18 years of age. Where parents object to LAIV on the grounds of its porcine gelatine content or where LAIV is unsuitable, children should be offered the injectable cell-based Quadrivalent Influenza Vaccine (QIVc), see Inactivated Influenza PGD.</p> <p>If the PGD is used for supply only for subsequent administration by an appropriately trained HCSW, the registered practitioner named in Section 7 of this PGD must supply the vaccine to the individual/carer. The HCSW cannot supply the medicine.</p> <p>Children with cochlear implants can be given LAIV safely although ideally not in the week prior to implant surgery or for two weeks afterwards, or if there is evidence of on-going cerebrospinal fluid leak.</p> <p>Exposure of healthcare professionals</p> <p>Very severely immunosuppressed individuals should not administer LAIV. Other healthcare workers who have less severe immunosuppression or are pregnant, should follow normal clinical practice to avoid inhaling the vaccine and ensure that they themselves are appropriately vaccinated.</p>
<p>Records</p> <p>Continued over page</p> <p>Records (continued)</p>	<p>Record:</p> <ul style="list-style-type: none"> • that valid informed consent was given • name of individual, address, date of birth and GP with whom the individual is registered (or record where an individual is not registered with a GP and that appropriate advice has been given) • clinical risk group indication for immunisation if applicable • name of immuniser • name and brand of vaccine • date of administration or supply • dose, form and route of administration of vaccine • quantity administered or supplied • batch number and expiry date • advice given; including advice given if excluded or declines immunisation • details of any adverse drug reactions and actions taken • whether supplied only or supplied and administered via PGD <p>Records should be signed and dated (or password-controlled immunisers record on e-records).</p> <p>All records should be clear, legible and contemporaneous.</p> <p>It is important that vaccinations given either at a general practice or elsewhere (for example, at schools or community pharmacies) are recorded on appropriate health records for the individual (using the appropriate clinical code). If given elsewhere, a record of vaccination should be returned to the individual's general practice to ensure a complete health record is held by the GP, allow clinical follow up and to avoid duplicate vaccination.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes in accordance with local policy.</p>

6. Key References

Key references	<p>LAIV</p> <ul style="list-style-type: none">• Immunisation Against Infectious Disease: The Green Book. Chapter 19, Updated 21 September 2022. Guidance: Influenza: the green book, chapter 19• Collection: Annual Flu Programme. Updated 25 May 2023 Annual flu programme - GOV.UK (www.gov.uk)• The national flu immunisation programme 2023 to 2024: supporting letter. Published 25 May 2023. National flu immunisation programme plan 2023 to 2024 - GOV.UK (www.gov.uk)• GP Contract: NHS England. england.nhs.uk/gp/investment/gp-contract/• Summary of Product Characteristics for Fluenz[®] Tetra. AstraZeneca UK Ltd. 22 March 2023. Fluenz Tetra nasal spray suspension Influenza vaccine (live attenuated, nasal) - Summary of Product Characteristics (SmPC) - (emc) (medicines.org.uk)• UKHSA Inactivated Influenza Vaccine PGD. government/publications/intramuscular-inactivated-influenza-vaccine-patient-group-direction-pgd-template• Flu Vaccinations: Supporting people with learning disabilities. 25 September 2018. government/publications/flu-vaccinations-for-people-with-learning-disabilities• Statement of amendment to the annual flu letter for 2023 to 2024 – 3 July 2023. www.gov.uk/government/publications/national-flu-immunisation-programme-plan/statement-of-amendment-to-the-annual-flu-letter-for-2023-to-2024-3-july-2023 <p>General</p> <ul style="list-style-type: none">• Health Technical Memorandum 07-01: Safe Management of Healthcare Waste. NHSE, 2022. NHS England » (HTM 07-01) Management and disposal of healthcare waste• Immunisation Against Infectious Disease: The Green Book. Chapter 2. Updated 18 June 2021. Consent: the green book, chapter 2 - GOV.UK (www.gov.uk)• National Minimum Standards and Core Curriculum for Immunisation Training. Published 7 February 2018. Immunisation training standards for healthcare practitioners - GOV.UK (www.gov.uk)• NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017. Overview Patient group directions Guidance NICE• NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. March 2017. Tools and resources Patient group directions Guidance NICE• Patient Group Directions: who can use them. Medicines and Healthcare products Regulatory Agency. 4 December 2017. Patient group directions: who can use them - GOV.UK (www.gov.uk)• UKHSA Immunisation Collection. www.gov.uk/government/collections/immunisation (www.gov.uk)• Vaccine Incident Guidance www.gov.uk/government/publications/vaccine-incident-guidance-responding-to-vaccine-errors
-----------------------	--

7. Practitioner authorisation sheet

LAIV PGD v13.00 Valid from: 1 September 2023 Expiry: 1 April 2024

Before signing this PGD, check that the document has had the necessary authorisations in section 2. Without these, this PGD is not lawfully valid.

Practitioner

By signing this PGD you are indicating that you agree to its contents and that you will work within it.

PGDs do not remove inherent professional obligations or accountability.

It is the responsibility of each professional to practise only within the bounds of their own competence and professional code of conduct.

I confirm that I have read and understood the content of this PGD and that I am willing and competent to work to it within my professional code of conduct.			
Name	Designation	Signature	Date

Authorising manager

I confirm that the practitioners named above have declared themselves suitably trained and competent to work under this PGD. I give authorisation on behalf of <u>insert name of organisation</u> for the above named health care professionals who have signed the PGD to work under it.			
Name	Designation	Signature	Date

Note to authorising manager

Score through unused rows in the list of practitioners to prevent practitioner additions post managerial authorisation.

This authorisation sheet should be retained to serve as a record of those practitioners authorised to work under this PGD.